

Literatur

Teil I Epidemiologie, Richtlinien und Organisation von Neugeborenen-Notarzdienst, Neonatologie und Geburtshilfe

- European Network for Perinatal Transport. Maternal and neonatal transport in Europe. Report of the European Network for Perinatal Transport (EUROPET). *Prenat Neonatal Med* 1999.
- Göpel W, Kribs A, Ziegler A et al. Avoidance of mechanical ventilation by surfactant treatment of spontaneously breathing preterm infants (AMV): an open-label, randomised, controlled trial. *Lancet* 2011; 378: 1627–34.
- Hansmann G. Neonatal resuscitation on air: it is time to turn down the oxygen tanks [corrected]. *Lancet* 2004; 364: 1293–4. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
- Hazinski MF, Nolan JP, Billi JE et al. Part 1: Executive summary: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation* 2010; 16 Suppl 2: S250–75.
- Hoehn T, Hansmann G, Bühner C et al. Therapeutic hypothermia in neonates. Review of current clinical data, ILCOR recommendations and suggestions for implementation in neonatal intensive care units. *Resuscitation* 2008; 1: 7–12. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
- Kribs A, Pillekamp F, Hunseler et al. Early administration of surfactant in spontaneous breathing with nCPAP: feasibility and outcome in extremely premature infants (postmenstrual age ≤ 27 weeks). *Paediatr Anaesth* 2007; 4: 364–9.
- Lucey JF. Why we should regionalize perinatal care. *Pediatrics* 1973; 4: 488–91.
- McNamara PJ, Mak W, Whyte HE. Dedicated neonatal retrieval teams improve delivery room resuscitation of outborn premature infants. *J Perinatol* 2005; 5: 309–14.
- Perlman JM, Wyllie J, Kattwinkel J et al. Part 11: neonatal resuscitation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation* 2010; 122 Suppl 2: S516–38. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
- Phibbs CS, Baker LC, Caughey AB et al. Level and volume of neonatal intensive care and mortality in very-low-birth-weight infants. *N Engl J Med* 2007; 21: 2165–75.
- Poets CF. What are the main research findings during the last 5 years that have changed my approach to clinical practice? *Arch Dis Child Fetal Neonatal Ed* 2013; 98: F89–92.
- Roehr CC, Hansmann G, Hoehn T, Bühner C. The 2010 Guidelines on Neonatal Resuscitation (AHA, ERC, ILCOR): similarities and differences – what progress has been made since 2005? *Klin Padiatr* 2011; 5: 299–307. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
- Saugstad OD. Resuscitation of newborn infants: from oxygen to room air. *Lancet* 2010; 376: 1970–1. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
- Sayre MR, O'Connor RE, Atkins DL et al. Part 2: evidence evaluation and management of potential or perceived conflicts of interest: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010; 18 Suppl 3: S657–64.
- Stark AR. Levels of neonatal care. *Pediatrics* 2004; 5: 1341–7.
- Van Reempts P, Gortner L, Milligan D et al. Characteristics of neonatal units that care for very preterm infants in Europe: results from the MOSAIC study. *Pediatrics* 2007; 4: e815–25.
- Wyllie J, Perlman JM, Kattwinkel J et al. Part 11: Neonatal resuscitation: 2010 International Con-

sensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Resuscitation 2010; 81 Suppl 1: e260–87. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in Circulation and Resuscitation.

18. Zeitlin J, Papiernik E, Breart G. Regionalization of perinatal care in Europe. *Semin Neonatol* 2004; 2: 99–110.

Teil II Basics für die Erstversorgung und Reanimation Neugeborener

1. Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000; 18: 1301–8.
2. Akisu M, Huseyinov A, Yalaz M et al. Selective head cooling with hypothermia suppresses the generation of platelet-activating factor in cerebrospinal fluid of newborn infants with perinatal asphyxia. *Prostaglandins Leukot Essent Fatty Acids* 2003; 1: 45–50.
3. Alderson PJ, Burrows FA, Stemp LI, Holtby HM. Use of ultrasound to evaluate internal jugular vein anatomy and to facilitate central venous cannulation in paediatric patients. *Br J Anaesth* 1993; 2: 145–8.
4. Allen HD, Driscoll DJ, Shaddy RE, Feltes TF. Moss and Adams' heart disease in infants, children, and adolescents: including the fetus and young adult. Philadelphia: Lippincott Williams & Wilkins 2012.
5. Aly H, Badawy M, El-Kholy A et al. Randomized, controlled trial on tracheal colonization of ventilated infants: can gravity prevent ventilator-associated pneumonia? *Pediatrics* 2008; 4: 770–4.
6. Ammari AN, Schulze KF. Uses and abuses of sodium bicarbonate in the neonatal intensive care unit. *Curr Opin Pediatr* 2002; 2: 151–6.
7. Anderson J, Leonard D, Braner DA et al. Videos in clinical medicine. Umbilical vascular catheterization. *N Engl J Med* 2008; 15: e18.
8. Andersson O, Hellstrom-Westas L, Andersson D, Domellof M. Effect of delayed versus early umbilical cord clamping on neonatal outcomes and iron status at 4 months: a randomised controlled trial. *BMJ* 2011; 343: d7157.
9. Extracorporeal life support organisation (ELSO), Ann Arbor. ELSO registry report (www.else.org/Registry.aspx).
10. Arai T, Yamashita M. Central venous catheterization in infants and children – small caliber audio-Doppler probe versus ultrasound scanner. *Paediatr Anaesth* 2005; 10: 858–61.
11. Arnold JH, Hanson JH, Toro-Figuero LO et al. Prospective, randomized comparison of high-frequency oscillatory ventilation and conventional mechanical ventilation in pediatric respiratory failure. *Crit Care Med* 1994; 10: 1530–9.
12. Askie LM, Henderson-Smart DJ, Irwig L, Simpson JM. Oxygen-saturation targets and outcomes in extremely preterm infants. *N Engl J Med* 2003; 10: 959–67.
13. Auten RL, Vozzelli M, Clark RH. Volutrauma. What is it, and how do we avoid it? *Clin Perinatol* 2001; 3: 505–15.
14. Azzopardi DV, Strohm B, Edwards AD et al. Moderate hypothermia to treat perinatal asphyxial encephalopathy. *N Engl J Med* 2009; 14: 1349–58.
15. Bahrami KR, Van Meurs KP. ECMO for neonatal respiratory failure. *Semin Perinatol* 2005; 1: 15–23.
16. Baker CF, Barks JD, Engmann C et al. Hydrocortisone administration for the treatment of refractory hypotension in critically ill newborns. *J Perinatol* 2008; 6: 412–9.
17. Baker PA, Brunette KE, Byrnes CA, Thompson JM. A prospective randomized trial comparing supraglottic airways for flexible bronchoscopy in children. *Paediatr Anaesth* 2010; 9: 831–8.
18. Ballard RA, Truog WE, Cnaan A et al. Inhaled nitric oxide in preterm infants undergoing mechanical ventilation. *N Engl J Med* 2006; 4: 343–53.
19. Barrington KJ, Byrne PJ. Premedication for neonatal intubation. *Am J Perinatol* 1998; 4: 213–6.
20. Barrington KJ. Umbilical artery catheters in the newborn: effects of position of the catheter tip. *Cochrane Database Syst Rev* 2000; 2: CD000505.
21. Barrington KJ, Finer NN. Inhaled nitric oxide for respiratory failure in preterm infants. *Cochrane Database Syst Rev* 2010; 10: CD000509.
22. Batton B, Batton D, Riggs T. Blood pressure during the first 7 days in premature infants born

- at postmenstrual age 23 to 25 weeks. *Am J Perinatol* 2007; 2: 107–15.
23. Beasley R, McNaughton A, Robinson G. New look at the oxyhaemoglobin dissociation curve. *Lancet* 2006; 367: 1124–6.
 24. Beck J, Campoccia F, Allo JC et al. Improved synchrony and respiratory unloading by neurally adjusted ventilatory assist (NAVA) in lung-injured rabbits. *Pediatric research* 2007; 3: 289–94.
 25. Beck J, Reilly M, Grasselli G et al. Patient-ventilator interaction during neurally adjusted ventilatory assist in low birth weight infants. *Pediatr Res* 2009; 6: 663–8.
 26. Bellomo R, Chapman M, Finfer S et al. Low-dose dopamine in patients with early renal dysfunction: a placebo-controlled randomised trial. Australian and New Zealand Intensive Care Society (ANZICS) Clinical Trials Group. *Lancet* 2000; 356: 2139–43.
 27. Berger TM, Bernet V, El Alama S et al. Perinatal care at the limit of viability between 22 and 26 completed weeks of gestation in Switzerland. 2011 revision of the Swiss recommendations. *Swiss Med Wkly* 2011; 141: w13280.
 28. Bernstein G, Knodel E, Heldt GP. Airway leak size in neonates and autocycling of three flow-triggered ventilators. *Crit Care Med* 1995; 10: 1739–44.
 29. Bhat MA, Shah ZA, Makhdoomi MS, Mufti MH. Theophylline for renal function in term neonates with perinatal asphyxia: a randomized, placebo-controlled trial. *J Pediatr* 2006; 2: 180–4.
 30. Bhuta T, Henderson-Smart DJ. Elective high-frequency oscillatory ventilation versus conventional ventilation in preterm infants with pulmonary dysfunction: systematic review and meta-analyses. *Pediatrics* 1997; 5: E6.
 31. Bittigau P, Sifringer M, Genz K et al. Antiepileptic drugs and apoptotic neurodegeneration in the developing brain. *Proc Natl Acad Sci U S A* 2002; 23: 15089–94.
 32. Blennow M, Jonsson B, Dahlstrom A et al. Lung function in premature infants can be improved. Surfactant therapy and CPAP reduce the need of respiratory support. *Lakartidningen* 1999; 13: 1571–6.
 33. Boloker J, Bateman DA, Wung JT, Stolar CJ. Congenital diaphragmatic hernia in 120 infants treated consecutively with permissive hypercapnea/spontaneous respiration/elective repair. *J Pediatr Surg* 2002; 3: 357–66.
 34. Bonifacio SL, Glass HC, Peloquin S, Ferriero DM. A new neurological focus in neonatal intensive care. *Nat Rev Neurol* 2011; 9: 485–94.
 35. Bouissou A, Rakza T, Klosowski S et al. Hypotension in preterm infants with significant patent ductus arteriosus: effects of dopamine. *J Pediatr* 2008; 6: 790–4.
 36. Bühner C, Bahr S, Siebert J et al. Use of 2% 2-phenoxyethanol and 0.1% octenidine as antiseptic in premature newborn infants of 23–26 weeks gestation. *J Hosp Infect* 2002; 4: 305–7.
 37. Cacciari A, Ruggeri G, Mordenti M et al. High-frequency oscillatory ventilation versus conventional mechanical ventilation in congenital diaphragmatic hernia. *Eur J Pediatr Surg* 2001; 1: 3–7.
 38. Carlo WA. Should nasal high-frequency ventilation be used in preterm infants? *Acta paediatrica* 2008; 11: 1484–5.
 39. Carlo WA, Finer NN, Walsh MC et al. Target ranges of oxygen saturation in extremely preterm infants. *N Engl J Med* 2010; 21: 1959–69.
 40. Carlton DP, Cummings JJ, Scheerer RG et al. Lung overexpansion increases pulmonary microvascular protein permeability in young lambs. *J Appl Physiol* 1990; 2: 577–83.
 41. Carlton DP, Cho SC, Davis P, Bland RD. Inflation pressure and lung vascular injury in preterm lambs. *Chest* 1994; 3 Suppl: 115–6S.
 42. Carter BS, Haverkamp AD, Merenstein GB. The definition of acute perinatal asphyxia. *Clin Perinatol* 1993; 2: 287–304.
 43. Casey BM, McIntire DD, Leveno KJ. The continuing value of the Apgar score for the assessment of newborn infants. *N Engl J Med* 2001; 7: 467–71.
 44. Cetta F, Graham LC, Eidem BW. Gaining vascular access in pediatric patients: use of the P.D. access Doppler needle. *Catheter Cardiovasc Interv* 2000; 1: 61–4.
 45. Clark RH, Gerstmann DR, Jobe AH et al. Lung injury in neonates: causes, strategies for prevention, and long-term consequences. *J Pediatr* 2001; 4: 478–86.
 46. Clark RH. The epidemiology of respiratory failure in neonates born at an estimated gestational age of 34 weeks or more. *J Perinatol* 2005; 4: 251–7.
 47. Cockburn F, Cooke R, Gamsu H. The CRIB (clinical risk index for babies) score: a tool for assessing initial neonatal risk and comparing

- performance of neonatal intensive care units. The International Neonatal Network. *Lancet* 1993; 342: 193–8.
48. Collins MP, Lorenz JM, Jetton JR, Paneth N. Hypocapnia and other ventilation-related risk factors for cerebral palsy in low birth weight infants. *Pediatr Res* 2001; 6: 712–9.
 49. Cools F, Henderson-Smart DJ, Offringa M, Askie LM. Elective high frequency oscillatory ventilation versus conventional ventilation for acute pulmonary dysfunction in preterm infants. *Cochrane Database Syst Rev* 2009; 3: CD000104.
 50. Cooper DS, Jacobs JP, Moore L et al. Cardiac extracorporeal life support: state of the art in 2007. *Cardiol Young* 2007; 17 Suppl 2: 104–15.
 51. Cotten CM. Air Leak Syndromes. In: Spitzer A (Hrsg). *Intensive Care of the Fetus and Neonate*. Philadelphia: Elsevier Mosby 2005.
 52. Courtney SE, Durand DJ, Asselin JM et al. High-frequency oscillatory ventilation versus conventional mechanical ventilation for very-low-birth-weight infants. *N Engl J Med* 2002; 9: 643–52.
 53. Courtney SE, Barrington KJ. Continuous positive airway pressure and noninvasive ventilation. *Clin Perinatol* 2007; 1: 73–92.
 54. Courtney SE, Barrington KJ. Continuous positive airway pressure and noninvasive ventilation. *Clin Perinatol* 2007; 1: 73–92.
 55. Dani C, Pratesi S, Migliori C, Bertini G. High flow nasal cannula therapy as respiratory support in the preterm infant. *Pediatric pulmonology* 2009; 7: 629–34.
 56. Dargaville PA. Innovation in surfactant therapy I: surfactant lavage and surfactant administration by fluid bolus using minimally invasive techniques. *Neonatology* 2012; 4: 326–36.
 57. Davis PG, Henderson-Smart DJ. Nasal continuous positive airways pressure immediately after extubation for preventing morbidity in preterm infants. *Cochrane Database Syst Rev* 2003; 2: CD000143.
 58. Davis PG, Tan A, O'Donnell CP, Schulze A. Resuscitation of newborn infants with 100% oxygen or air: a systematic review and meta-analysis. *Lancet* 2004; 364: 1329–33.
 59. Davis PG, Morley CJ, Owen LS. Non-invasive respiratory support of preterm neonates with respiratory distress: continuous positive airway pressure and nasal intermittent positive pressure ventilation. *Semin Fetal Neonatal Med* 2009; 1: 14–20.
 60. Dawes GS. Birth asphyxia, resuscitation, and brain damage (chap 12). *Foetal and neonatal physiology*. Chicago: Year Book Publisher 1968; 141–59.
 61. Day RW, Lynch JM, White KS, Ward RM. Acute response to inhaled nitric oxide in newborns with respiratory failure and pulmonary hypertension. *Pediatrics* 1996; 4 Pt 1: 698–705.
 62. Dempsey EM, Al Hazzani F, Faucher D, Barrington KJ. Facilitation of neonatal endotracheal intubation with mivacurium and fentanyl in the neonatal intensive care unit. *Arch Dis Child Fetal Neonatal Ed* 2006; 4: F279–82.
 63. Dempsey EM, Barrington KJ. Treating hypotension in the preterm infant: when and with what: a critical and systematic review. *J Perinatol* 2007; 8: 469–78.
 64. Dempsey EM, Barrington KJ. Evaluation and treatment of hypotension in the preterm infant. *Clin Perinatol* 2009; 1: 75–85.
 65. Dempsey EM, Al Hazzani F, Barrington KJ. Permissive hypotension in the extremely low birthweight infant with signs of good perfusion. *Arch Dis Child Fetal Neonatal Ed* 2009; 4: F241–4.
 66. Dettenkofer M, Wenzler S, Amthor S et al. Does disinfection of environmental surfaces influence nosocomial infection rates? A systematic review. *Am J Infect Control* 2004; 2: 84–9.
 67. Deutsche Gesellschaft für Gynäkologie und Geburtshilfe, Deutsche Gesellschaft für Kinder- und Jugendmedizin, Deutsche Gesellschaft für Perinatale Medizin, Akademie für Ethik in der Medizin, Gesellschaft für Neonatologie und Pädiatrische Intensivmedizin, Deutscher Hebammenverband, Bundesverband »Das frühgeborene Kind«. *Frühgeborene an der Grenze der Lebensfähigkeit (AWMF-Register Nr. 024/019, bitte Update von 04/2014 lesen unter www.awmf.org)*.
 68. Deutsche Gesellschaft für Gynäkologie und Geburtshilfe (DGGG), Deutsche Gesellschaft für Kinder- und Jugendmedizin (DGKJ), Deutsche Gesellschaft für Perinatale Medizin (DGPM), Deutscher Hebammenverband, Gesellschaft für Neonatologie und Pädiatrische Intensivmedizin (GNPI). *Betreuung von gesunden reifen Neugeborenen in der Geburtsklinik (AWMF-Register Nr. 024/005; www.awmf.org)*.

69. DiBlasi RM. Neonatal noninvasive ventilation techniques: do we really need to intubate? *Respir Care* 2011; 9: 1273–94; discussion 95–7.
70. Dixon H, Hawkins K, Stephenson T. Comparison of albumin versus bicarbonate treatment for neonatal metabolic acidosis. *Eur J Pediatr* 1999; 5: 414–5.
71. Duffett M, Choong K, Ng V et al. Surfactant therapy for acute respiratory failure in children: a systematic review and meta-analysis. *Crit Care* 2007; 3: R66.
72. Duman N, Tuzun F, Sutcuoglu S et al. Impact of volume guarantee on synchronized ventilation in preterm infants: a randomized controlled trial. *Intensive Care Med* 2012; 8: 1358–64.
73. Edwards AD, Brocklehurst P, Gunn AJ et al. Neurological outcomes at 18 months of age after moderate hypothermia for perinatal hypoxic ischaemic encephalopathy: synthesis and meta-analysis of trial data. *BMJ* 2010; 340: c363.
74. Eicher DJ, Wagner CL, Katikaneni LP et al. Moderate hypothermia in neonatal encephalopathy: efficacy outcomes. *Pediatr Neurol* 2005; 1: 11–7.
75. Engle WA. Intraosseous access for administration of medications in neonates. *Clin Perinatol* 2006; 1: 161–8, ix.
76. European Resuscitation Council. Newborn Life Support (Poster). Oktober 2010. www.erc.edu/index.php/view_category/en/cid=12.
77. Ewer AK, Middleton LJ, Furmston AT et al. (PulseOx Study Group). Pulse oximetry screening for congenital heart defects in newborn infants (PulseOx): a test accuracy study. *Lancet* 2011; 378: 785–94.
78. Fabres J, Carlo WA, Phillips V et al. Both extremes of arterial carbon dioxide pressure and the magnitude of fluctuations in arterial carbon dioxide pressure are associated with severe intraventricular hemorrhage in preterm infants. *Pediatrics* 2007; 2: 299–305.
79. Falck AJ, Escobedo MB, Baillargeon JG et al. Proficiency of pediatric residents in performing neonatal endotracheal intubation. *Pediatrics* 2003; 6 Pt 1: 1242–7.
80. Fanconi G, Lemburg P. Volumenersatz beim Früh- und Neugeborenen. In: Schöber JG, Lemburg P (eds). *Erstversorgung von Risikoneugeborenen im Kreißsaal und auf dem Transport*. München: Alete Wissenschaftlicher Dienst 1999; 30–3.
81. Filippi L, Pezzati M, Poggi C et al. Dopamine versus dobutamine in very low birthweight infants: endocrine effects. *Arch Dis Child Fetal Neonatal Ed* 2007; 5: F367–71.
82. Finer NN, Carlo WA, Duara S et al. (National Institute of Child Health, Human Development Neonatal Research Network). Delivery room continuous positive airway pressure/positive end-expiratory pressure in extremely low birth weight infants: a feasibility trial. *Pediatrics* 2004; 3: 651–7.
83. Finer NN, Rich WD. Neonatal resuscitation: raising the bar. *Curr Opin Pediatr* 2004; 2: 157–62.
84. Finer NN, Carlo WA, Walsh MC et al. Early CPAP versus surfactant in extremely preterm infants. *N Engl J Med* 2010; 21: 1970–9.
85. Fiorito BA, Mirza F, Doran TM et al. Intraosseous access in the setting of pediatric critical care transport. *Pediatr Crit Care Med* 2005; 1: 50–3.
86. Fischer HS, Roehr CC, Proquitte H et al. Is volume and leak monitoring feasible during nasopharyngeal continuous positive airway pressure in neonates? *Intensive Care Med* 2009; 11: 1934–41.
87. Fujiwara T, Maeta H, Chida S et al. Artificial surfactant therapy in hyaline-membrane disease. *Lancet* 1980; 1: 55–9.
88. Gemeinsamer Bundesausschuss (G-BA). Beschluss des Gemeinsamen Bundesausschusses über eine Beauftragung des Instituts für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG): Bewertung eines Screenings auf schwere congenitale Herzfehler mittels Pulsoxymetrie. www.g-ba.de/downloads/39-261-1744/2013-06-20_IQWiG_Beauftragung-Pulsoxymetrie.pdf.
89. Gesellschaft für Neonatologie und Pädiatrische Intensivmedizin, Deutsche Gesellschaft für Perinatale Medizin, Deutsche Diabetes-Gesellschaft, Deutsche Gesellschaft für Kinder- und Jugendmedizin, Deutsche Gesellschaft für Gynäkologie und Geburtshilfe. *Betreuung von Neugeborenen diabetischer Mütter (AWMF-Register Nr. 024/006; www.awmf.org)*.
90. Ghanta S, Abdel-Latif ME, Lui K et al. Propofol compared with the morphine, atropine, and suxamethonium regimen as induction agents for neonatal endotracheal intubation: a randomized, controlled trial. *Pediatrics* 2007; 119: e1248–55.

91. Glass P, Wagner AE, Papero PH et al. Neurodevelopmental status at age five years of neonates treated with extracorporeal membrane oxygenation. *J Pediatr* 1995; 3: 447–57.
92. Gluckman PD, Wyatt JS, Azzopardi D et al. Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicentre randomised trial. *Lancet* 2005; 365: 663–70.
93. Goldmann K, Roettger C, Wulf H. The size 1(1/2) ProSeal laryngeal mask airway in infants: a randomized, crossover investigation with the Classic laryngeal mask airway. *Anesth Analg* 2006; 2: 405–10.
94. Gopel W, Kribs A, Ziegler A et al. Avoidance of mechanical ventilation by surfactant treatment of spontaneously breathing preterm infants (AMV): an open-label, randomised, controlled trial. *Lancet* 2011; 378: 1627–34.
95. Greenough A, Milner AD, Dimitriou G. Synchronized mechanical ventilation for respiratory support in newborn infants. *Cochrane Database Syst Rev* 2004; 4: CD000456.
96. Gunn AJ, Hoehn T, Hansmann G et al. Hypothermia, an evolving treatment for neonatal hypoxic ischemic encephalopathy. *Pediatrics* 2008; 121: 648–9.
97. Guntheroth W. *Physiology of the circulation; fetus, neonate and child*. Philadelphia: Harper & Row 1983.
98. Haas NA, Haas SA. Central venous catheter techniques in infants and children. *Curr Opin Anaesthesiol* 2003; 3: 291–303.
99. Haas NA. Clinical review: vascular access for fluid infusion in children. *Crit Care* 2004; 6: 478–84.
100. Hahn S, Choi HJ, Soll R, Dargaville PA. Lung lavage for meconium aspiration syndrome in newborn infants. *Cochrane Database Syst Rev* 2013; 4: CD003486.
101. Halamek LP, Morley C. Continuous positive airway pressure during neonatal resuscitation. *Clin Perinatol* 2006; 1: 83–98, vii.
102. Hall RW, Kronsberg SS, Barton BA et al. Morphine, hypotension, and adverse outcomes among preterm neonates: who's to blame? Secondary results from the NEOPAIN trial. *Pediatrics* 2005; 5: 1351–9.
103. Hallberg B, Olson L, Bartocci M et al. Passive induction of hypothermia during transport of asphyxiated infants: a risk of excessive cooling. *Acta Paediatr* 2009; 6: 942–6.
104. Halliday HL. Endotracheal intubation at birth for preventing morbidity and mortality in vigorous, meconium-stained infants born at term. *Cochrane Database Syst Rev* 2001; 1: CD000500.
105. Halliday HL, Ehrenkranz RA, Doyle LW. Early (<8 days) postnatal corticosteroids for preventing chronic lung disease in preterm infants. *Cochrane Database Syst Rev* 2010; 1: CD001146.
106. Hansmann G. Neonatal resuscitation on air: it is time to turn down the oxygen tanks [corrected]. *Lancet* 2004; 364: 1293–4.
107. Hansmann G (ed). *Neonatal Emergencies*. Cambridge University Press 2009.
108. Heine VM, Rowitch DH. Hedgehog signaling has a protective effect in glucocorticoid-induced mouse neonatal brain injury through an 11betaHSD2-dependent mechanism. *J Clin Invest* 2009; 2: 267–77.
109. Henderson-Smart DJ, De Paoli AG, Clark RH, Bhuta T. High frequency oscillatory ventilation versus conventional ventilation for infants with severe pulmonary dysfunction born at or near term. *Cochrane Database Syst Rev* 2009; 3: CD002974.
110. Herrera CM, Gerhardt T, Claire N et al. Effects of volume-guaranteed synchronized intermittent mandatory ventilation in preterm infants recovering from respiratory failure. *Pediatrics* 2002; 3: 529–33.
111. Herting E, Gan X, Rauprich P et al. Combined treatment with surfactant and specific immunoglobulin reduces bacterial proliferation in experimental neonatal group B streptococcal pneumonia. *Am J Respir Crit Care Med* 1999; 6: 1862–7.
112. Herting E, Schiffmann H, Roth C, Johansson J. Surfactant lavage demonstrates protein fibrils in a neonate with congenital surfactant protein b deficiency. *Am J Respir Crit Care Med* 2002; 9: 1292–4.
113. Higgs ZC, Macafee DA, Braithwaite BD et al. The Seldinger technique: 50 years on. *Lancet* 2005; 366: 1407–9.
114. Hillewig E, Aghayev E, Jackowski C et al. Gas embolism following intraosseous medication application proven by post-mortem multislice computed tomography and autopsy. *Resuscitation* 2007; 1: 149–53.
115. Hines MH. ECMO and congenital heart disease. *Semin Perinatol* 2005; 1: 34–9.

116. Hoehn T, Krause MF, Bühner C. Meta-analysis of inhaled nitric oxide in premature infants: an update. *Klin Padiatr* 2006; 2: 57–61.
117. Hoehn T, Hansmann G, Bühner C et al. Therapeutic hypothermia in neonates. Review of current clinical data, ILCOR recommendations and suggestions for implementation in neonatal intensive care units. *Resuscitation* 2008; 1: 7–12.
118. Hoffman TM, Wernovsky G, Atz AM et al. Efficacy and safety of milrinone in preventing low cardiac output syndrome in infants and children after corrective surgery for congenital heart disease. *Circulation* 2003; 7: 996–1002.
119. Huang QW, Sun B, Gao F et al. Effects of inhaled nitric oxide and high-frequency ventilation in rabbits with meconium aspiration. *Biol Neonate* 1999; 6: 374–82.
120. Hutton EK, Hassan ES. Late vs early clamping of the umbilical cord in full-term neonates: systematic review and meta-analysis of controlled trials. *JAMA* 2007; 11: 1241–52.
121. ILCOR. International consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. Part 7. Neonatal Resuscitation. *Resuscitation* 2005; 67: 293–303.
122. Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG). Screening auf kritische angeborene Herzfehler mittels Pulsoxymetrie bei Neugeborenen. [S13-01]. www.iqwig.de/de/projekte-ergebnisse/projekte/nichtmedikamentose-verfahren/s13-01-screening-auf-kritische-angeborene-herzfehler-mittels-pulsoxymetrie-bei-neugeborenen.3681.html.
123. Jacobs SE, Morley CJ, Inder TE et al. Whole-body hypothermia for term and near-term newborns with hypoxic-ischemic encephalopathy: a randomized controlled trial. *Arch Pediatr Adolesc Med* 2011; 165: 692–700.
124. Jobe AH, Bancalari E. Bronchopulmonary dysplasia. *Am J Respir Crit Care Med* 2001; 7: 1723–9.
125. Jobe AH, Kramer BW, Moss TJ et al. Decreased indicators of lung injury with continuous positive expiratory pressure in preterm lambs. *Pediatr Res* 2002; 3: 387–92.
126. Joshi G, Tobias JD. A five-year experience with the use of BiPAP in a pediatric intensive care unit population. *J Intensive Care Med* 2007; 1: 38–43.
127. Kabra NS, Kumar M, Shah SS. Multiple versus single lumen umbilical venous catheters for newborn infants. *Cochrane Database Syst Rev* 2005; 3: CD004498.
128. Kattwinkel J, Bloom BT, Delmore P et al. High-versus low-threshold surfactant retreatment for neonatal respiratory distress syndrome. *Pediatrics* 2000; 2 Pt 1: 282–8.
129. Kattwinkel J. Surfactant lavage for meconium aspiration: a word of caution. *Pediatrics* 2002; 6: 1167–8.
130. Kattwinkel J. Neonatal Resuscitation. Elk Grove Village, IL: American Academy of Pediatrics and American Heart Association 2006.
131. Kattwinkel J, Perlman JM, Aziz K et al. Part 15: neonatal resuscitation: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010; 18 Suppl 3: S909–19.
132. Kattwinkel J. Textbook of neonatal resuscitation: American Academy of Pediatrics 2011.
133. Kellum JA, J MD. Use of dopamine in acute renal failure: a meta-analysis. *Crit Care Med* 2001; 8: 1526–31.
134. Kinsella JP, Cutter GR, Walsh WF et al. Early inhaled nitric oxide therapy in premature newborns with respiratory failure. *N Engl J Med* 2006; 4: 354–64.
135. Kliegman R. Fetal and neonatal medicine. In: Behrman R, Kliegman R (eds). *Nelson Essentials of Pediatrics*. Philadelphia: W.R. Saunders Company 2002; 179–249.
- 135a. Kommission für Krankenhaushygiene und Infektionsprävention (KRINKO). Empfehlung zur Prävention nosokomialer Infektionen bei neonatologischen Intensivpflegepatienten mit einem Geburtsgewicht unter 1 500 g. *Bundesgesundheitsbl – Gesundheitsforsch – Gesundheitsschutz* 2007; 50: 1265–303.
- 135b. Kommission für Krankenhaushygiene und Infektionsprävention (KRINKO). Ergänzende Empfehlung (2011) zur »Prävention nosokomialer Infektionen bei neonatologischen Intensivpflegepatienten mit einem Geburtsgewicht unter 1 500 g« (2007). *Epidemiologisches Bulletin* 2012; 2: 13–8.
136. Kribs A. Is it safer to intubate premature infants in the delivery room? *Pediatrics* 2006; 5: 1858–9; author reply 9.
137. Kribs A, Pillekamp F, Hunseler C et al. Early administration of surfactant in spontaneous

- breathing with nCPAP: feasibility and outcome in extremely premature infants (postmenstrual age $</=27$ weeks). *Paediatr Anaesth* 2007; 4: 364–9.
138. Kribs A, Hartel C, Kattner E et al. Surfactant without intubation in preterm infants with respiratory distress: first multi-center data. *Klin Padiatr* 2010; 1: 13–7.
 139. Kugelman A, Gangitano E, Taschuk R et al. Extracorporeal membrane oxygenation in infants with meconium aspiration syndrome: a decade of experience with venovenous ECMO. *J Pediatr Surg* 2005; 7: 1082–9.
 140. Lake W, Emmerson AJ. Use of a butterfly as an intraosseous needle in an oedematous preterm infant. *Arch Dis Child Fetal Neonatal Ed* 2003; 5: F409.
 141. Lally KP, Lally PA, Van Meurs KP et al. Treatment evolution in high-risk congenital diaphragmatic hernia: ten years' experience with diaphragmatic agenesis. *Ann Surg* 2006; 4: 505–13.
 142. Latto IP, Rosen M. Percutaneous central venous and arterial catheterisation. London, New York: Saunders 2000.
 143. Lemburg P. Erstversorgung von Risikoneugeborenen im Kreißsaal und auf dem Transport. *Alete Wissenschaftlicher Dienst* 1999.
 144. Lemburg P. Neonatale Hypoxie (Asphyxie), Reanimation und Folgen. *Schöber JG* 1999.
 145. Leone TA, Lange A, Rich W, Finer NN. Disposable colorimetric carbon dioxide detector use as an indicator of a patent airway during noninvasive mask ventilation. *Pediatrics* 2006.
 146. Leone TA, Rich W, Finer NN. A survey of delivery room resuscitation practices in the United States. *Pediatrics* 2006; 2: e164–75.
 147. Lin HC, Su BH, Lin TW et al. System-based strategy for the management of meconium aspiration syndrome: 198 consecutive cases observations. *Acta Paediatr Taiwan* 2005; 2: 67–71.
 148. Lin ZL, Yu HM, Lin J et al. Mild hypothermia via selective head cooling as neuroprotective therapy in term neonates with perinatal asphyxia: an experience from a single neonatal intensive care unit. *J Perinatol* 2006; 3: 180–4.
 149. Lindner W, Pohlandt F. Oxygenation and ventilation in spontaneously breathing very preterm infants with nasopharyngeal CPAP in the delivery room. *Acta Paediatr Scand* 2007; 1: 17–22.
 150. Locke RG, Wolfson MR, Shaffer TH et al. Inadvertent administration of positive end-distending pressure during nasal cannula flow. *Pediatrics* 1993; 1: 135–8.
 151. Logan JW, O'Shea TM, Allred EN et al. Early postnatal hypotension and developmental delay at 24 months of age among extremely low gestational age newborns. *Arch Dis Child Fetal Neonatal Ed* 2011; 5: F321–8.
 152. Lopez-Gil M, Mantilla I, Blanco T et al. The Size 1 ProSeal laryngeal mask airway in infants: a randomized, noncrossover study with the Classic laryngeal mask airway. *Paediatr Anaesth* 2012; 4: 365–70.
 153. Lundstrom KE, Pryds O, Greisen G. Oxygen at birth and prolonged cerebral vasoconstriction in preterm infants. *Arch Dis Child Fetal Neonatal Ed* 1995; 2: F81–6.
 154. Mahle W, Koppel R. Screening with pulse oximetry for congenital heart disease. *Lancet* 2011; 378: 749–50.
 155. Mahle WT, Martin GR, Beekman RH et al. Endorsement of Health and Human Services recommendation for pulse oximetry screening for critical congenital heart disease. *Pediatrics* 2012; 1: 190–2.
 156. Mahmoud RA, Roehr CC, Schmalisch G. Current methods of non-invasive ventilatory support for neonates. *Paediatr Respir Rev* 2011; 3: 196–205.
 157. Mahmoud RA, Proquitte H, Fawzy N et al. Tracheal tube airleak in clinical practice and impact on tidal volume measurement in ventilated neonates. *Pediatr Crit Care Med* 2011; 2: 197–202.
 158. Marik PE. Low-dose dopamine: a systematic review. *Intensive Care Med* 2002; 7: 877–83.
 159. Marlow N, Greenough A, Peacock JL et al. Randomised trial of high frequency oscillatory ventilation or conventional ventilation in babies of gestational age 28 weeks or less: respiratory and neurological outcomes at 2 years. *Arch Dis Child Fetal Neonatal Ed* 2006; 5: F320–6.
 160. McCallion N, Davis PG, Morley CJ. Volume-targeted versus pressure-limited ventilation in the neonate. *Cochrane Database Syst Rev* 2005; 3: CD003666.
 161. McDonald SJ, Middleton P. Effect of timing of umbilical cord clamping of term infants on maternal and neonatal outcomes. *Cochrane Database Syst Rev* 2008; 2: CD004074.
 162. McGuire W, Fowlie PW. Naloxone for narcotic-exposed newborn infants. *Cochrane Database Syst Rev* 2002; 4: CD003483.

163. McNally H, Bennett CC, Elbourne D, Field DJ, Group UKCET. United Kingdom collaborative randomized trial of neonatal extracorporeal membrane oxygenation: follow-up to age 7 years. *Pediatrics* 2006; 5: e845–54.
164. Mellon RD, Simone AF, Rappaport BA. Use of anesthetic agents in neonates and young children. *Anesth Analg* 2007; 3: 509–20.
165. Mercier JC, Hummler H, Durrmeyer X et al. (Group ES). Inhaled nitric oxide for prevention of bronchopulmonary dysplasia in premature babies (EUNO): a randomised controlled trial. *Lancet* 2010; 376: 346–54.
166. Micaglio M, Trevisanuto D, Doglioni N et al. The size 1 LMA-ProSeal: Comparison with the LMA-Classic during pressure controlled ventilation in a neonatal intubation manikin. *Resuscitation* 2007; 1: 124–7.
167. Migliori C, Motta M, Angeli A, Chirico G. Nasal bilevel vs. continuous positive airway pressure in preterm infants. *Pediatr Pulmonol* 2005; 5: 426–30.
168. Millar KJ, Thiagarajan RR, Laussen PC. Glucocorticoid therapy for hypotension in the cardiac intensive care unit. *Pediatr Cardiol* 2007; 3: 176–82.
169. Mizuno K, Takeuchi T, Itabashi K, Okuyama K. Efficacy of synchronized IMV on weaning neonates from the ventilator. *Acta Paediatr Jpn* 1994; 2: 162–6.
170. Morley CJ, Davis PG, Doyle LW et al. Nasal CPAP or intubation at birth for very preterm infants. *N Engl J Med* 2008; 7: 700–8.
171. Naik AS, Kallapur SG, Bachurski CJ et al. Effects of ventilation with different positive end-expiratory pressures on cytokine expression in the preterm lamb lung. *Am J Respir Crit Care Med* 2001; 3: 494–8.
172. Nelson KB, Emery ES, 3rd. Birth asphyxia and the neonatal brain: what do we know and when do we know it? *Clin Perinatol* 1993; 2: 327–44.
173. Nichols DG. *Critical heart disease in infants and children*. Philadelphia: Mosby 2006.
174. Niermeyer S, Kattwinkel J, Van Reempts P et al. International Guidelines for Neonatal Resuscitation: An excerpt from the Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care: International Consensus on Science. Contributors and Reviewers for the Neonatal Resuscitation Guidelines. *Pediatrics* 2000; 3: E29.
175. Niermeyer S, Vento M. Is 100% oxygen necessary for the resuscitation of newborn infants? *J Matern Fetal Neonatal Med* 2004; 2: 75–84.
176. Nolan JP, Soar J, Zideman DA et al. (Group ERCGW). European Resuscitation Council Guidelines for Resuscitation 2010 Section 1. Executive summary. *Resuscitation* 2010; 10: 1219–76. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
177. Nolan JP, Soar J, Perkins GD. Resuscitation 2010 in review. *Resuscitation* 2011; 2: 145–8.
178. Noori S, Seri I. Neonatal blood pressure support: the use of inotropes, lusitropes, and other vasopressor agents. *Clin Perinatol* 2012; 1: 221–38.
179. O'Donnell C, Davis P, Morley C. Positive end-expiratory pressure for resuscitation of newborn infants at birth. *Cochrane Database Syst Rev* 2004; 4: CD004341.
180. O'Donnell CP, Davis PG, Morley CJ. Positive pressure ventilation at neonatal resuscitation: review of equipment and international survey of practice. *Acta Paediatr Scand* 2004; 5: 583–8.
181. O'Donnell CP, Davis PG, Lau R et al. Neonatal resuscitation 2: an evaluation of manual ventilation devices and face masks. *Arch Dis Child Fetal Neonatal Ed* 2005; 5: F392–6.
182. O'Donnell CP, Kamlin CO, Davis PG, Morley CJ. Endotracheal intubation attempts during neonatal resuscitation: success rates, duration, and adverse effects. *Pediatrics* 2006; 1: e16–21.
183. O'Grady NP, Alexander M, Dellinger EP et al. Guidelines for the prevention of intravascular catheter-related infections. The Hospital Infection Control Practices Advisory Committee, Center for Disease Control and Prevention, u.s. *Pediatrics* 2002; 5: e51.
184. Obladen M (Hrsg). *Neugeborenenintensivpflege*. Berlin, Heidelberg: Springer 2002.
185. Okuyama H, Kubota A, Oue T et al. Inhaled nitric oxide with early surgery improves the outcome of antenatally diagnosed congenital diaphragmatic hernia. *J Pediatr Surg* 2002; 8: 1188–90.
186. Orkin SH, Nathan D, Ginsburg D. *Nathan and Oski's Hematology of Infancy and Childhood*: Saunders W B Co 2008.
187. Osborn DA, Paradis M, Evans N. The effect of inotropes on morbidity and mortality in

- preterm infants with low systemic or organ blood flow. *Cochrane Database Syst Rev* 2007; 1: CD005090.
188. Owen CJ, Wyllie JP. Determination of heart rate in babies at birth. *Resuscitation* 2004; 60: 213–7.
 189. Owen LS, Morley CJ, Davis PG. Neonatal nasal intermittent positive pressure ventilation: what do we know in 2007? *Archives of disease in childhood Fetal and neonatal edition* 2007; 5: F414–8.
 190. Owen LS, Morley CJ, Davis PG. Neonatal nasal intermittent positive pressure ventilation: a survey of practice in England. *Arch Dis Child Fetal Neonatal Ed* 2008; 93: F148–50.
 191. Paradisis M, Evans N, Kluckow M, Osborn D. Randomized trial of milrinone versus placebo for prevention of low systemic blood flow in very preterm infants. *J Pediatr* 2009; 2: 189–95.
 192. Park MK (ed). *Pediatric cardiology for practitioners*. St. Louis: Mosby 2014.
 193. Perlman JM, Wyllie J, Kattwinkel J et al. Part 11: neonatal resuscitation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation* 2010; 16 Suppl 2: S516–38.
 194. Pillow JJ. High-frequency oscillatory ventilation: mechanisms of gas exchange and lung mechanics. *Crit Care Med* 2005; 3 Suppl: S135–41.
 195. Pillow JJ, Minocchieri S. Innovation in surfactant therapy II: surfactant administration by aerosolization. *Neonatology* 2012; 4: 337–44.
 196. Poets CF. What are the main research findings during the last 5 years that have changed my approach to clinical practice? *Arch Dis Child Fetal Neonatal Ed* 2013; 98: F89–92.
 197. Polin RA, Carlo WA, Committee on F, Newborn, American Academy of P. Surfactant replacement therapy for preterm and term neonates with respiratory distress. *Pediatrics* 2014; 1: 156–63.
 198. Rabe H, Reynolds G, Diaz-Rossello J. A systematic review and meta-analysis of a brief delay in clamping the umbilical cord of preterm infants. *Neonatology* 2008; 2: 138–44.
 199. Ralston M, Hazinski MF, Zaritsky AL et al. *PALS Provider Manual*. Dallas: American Heart Association and American Academy of Pediatrics 2006.
 200. Rennie JM. *Robertson's Textbook of Neonatology*. Edinburgh: Churchill Livingstone 2005.
 201. Rennie JM. *Rennie & Robertson's Textbook of Neonatology*. Edinburgh: Churchill Livingstone 2012.
 202. Rennie JR, NRC. *Textbook of Neonatology*. Edinburgh: Churchill Livingstone 1999.
 203. Richmond S. *ILCOR and neonatal resuscitation* 2005. *Arch Dis Child Fetal Neonatal Ed* 2007; 3: F163–5.
 204. Richmond S, Wyllie J. *European Resuscitation Council Guidelines for Resuscitation 2010 Section 7. Resuscitation of babies at birth*. *Resuscitation* 2010; 10: 1389–99.
 205. Roberts KD, Leone TA, Edwards WH et al. Pre-medication for nonemergent neonatal intubations: a randomized, controlled trial comparing atropine and fentanyl to atropine, fentanyl, and mivacurium. *Pediatrics* 2006; 4: 1583–91.
 206. Robertson NJ, Nakakeeto M, Hagmann C et al. Therapeutic hypothermia for birth asphyxia in low-resource settings: a pilot randomised controlled trial. *Lancet* 2008; 372: 801–3.
 207. Roehr CC, Hansmann G, Hoehn T, Bührer C. The 2010 Guidelines on Neonatal Resuscitation (AHA, ERC, ILCOR): similarities and differences – what progress has been made since 2005? *Klin Padiatr* 2011; 5: 299–307.
 208. Rojas-Reyes MX, Morley CJ, Soll R. Prophylactic versus selective use of surfactant in preventing morbidity and mortality in preterm infants. *Cochrane Database Syst Rev* 2012; 3: CD000510.
 209. Roth P, Harris M, Vega-Rich C, Marro P. *Neonatology*. In: Polin RA DM (Hrsg). *Pediatric Secrets*. Philadelphia: Hanley & Belfus 2001; 409–65.
 210. Rudolph AM. *Congenital diseases of the heart: clinical-physiological considerations*. Hoboken: Wiley-Blackwell 2009.
 211. Rutherford M, Ramenghi LA, Edwards AD et al. Assessment of brain tissue injury after moderate hypothermia in neonates with hypoxic-ischaemic encephalopathy: a nested substudy of a randomised controlled trial. *Lancet Neurol* 2010; 1: 39–45.
 212. Rutherford MA, Azzopardi D, Whitelaw A et al. Mild hypothermia and the distribution of cerebral lesions in neonates with hypoxic-ischemic encephalopathy. *Pediatrics* 2005; 4: 1001–6.
 213. Saarenmaa E, Huttunen P, Leppaluoto J et al. Advantages of fentanyl over morphine in analgesia for ventilated newborn infants after birth: a randomized trial. *J Pediatr* 1999; 2: 144–50.

214. Sarnat HB, Sarnat MS. Neonatal encephalopathy following fetal distress. A clinical and electroencephalographic study. *Arch Neurol* 1976; 10: 696–705.
215. Saugstad OD, Rootwelt T, Aalen O. Resuscitation of asphyxiated newborn infants with room air or oxygen: an international controlled trial: the Resair 2 study. *Pediatrics* 1998; 1: e1.
216. Saugstad OD, Ramji S, Vento M. Resuscitation of depressed newborn infants with ambient air or pure oxygen: a meta-analysis. *Biol Neonate* 2005; 1: 27–34.
217. Saugstad OD, Ramji S, Vento M. Resuscitation of depressed newborn infants with ambient air or pure oxygen: a meta-analysis. *Biol Neonate* 2005; 1: 27–34.
218. Saugstad OD. Oxygen saturations immediately after birth. *J Pediatr* 2006; 5: 569–70.
219. Saugstad OD, Ramji S, Soll RF, Vento M. Resuscitation of newborn infants with 21 % or 100 % oxygen: an updated systematic review and meta-analysis. *Neonatology* 2008; 3: 176–82.
220. Saugstad OD. Resuscitation of newborn infants: from oxygen to room air. *Lancet* 2010; 376: 1970–1.
221. Schaible T, Hermle D, Loersch F et al. A 20-year experience on neonatal extracorporeal membrane oxygenation in a referral center. *Intensive Care Med* 2010; 7: 1229–34.
222. Schmolzer GM, Kamlin CO, Dawson JA et al. Tidal volume delivery during surfactant administration in the delivery room. *Intensive Care Med* 2011; 11: 1833–9.
223. Segerer H, van Gelder W, Angenent FW et al. Pulmonary distribution and efficacy of exogenous surfactant in lung-lavaged rabbits are influenced by the instillation technique. *Pediatr Res* 1993; 4: 490–4.
224. Shankaran S, Laptook A, Wright LL et al. Whole-body hypothermia for neonatal encephalopathy: animal observations as a basis for a randomized, controlled pilot study in term infants. *Pediatrics* 2002; 2 Pt 1: 377–85.
225. Shankaran S, Laptook AR, Ehrenkranz RA et al. (National Institute of Child Health, Human Development Neonatal Research Network). Whole-body hypothermia for neonates with hypoxic-ischemic encephalopathy. *N Engl J Med* 2005; 15: 1574–84.
226. Shankaran S, Laptook AR, Ehrenkranz RA et al. Whole-body hypothermia for neonates with hypoxic-ischemic encephalopathy. *N Engl J Med* 2005; 15: 1574–84.
227. Shem S. House of God. New York: Dell Publishing 1978.
228. Simbruner G, Mittal RA, Rohlmann F, Muche R. Systemic hypothermia after neonatal encephalopathy: outcomes of neo.nEURO.network RCT. *Pediatrics* 2010; 4: e771–8.
229. Singh J, Sinha SK, Clarke P et al. Mechanical ventilation of very low birth weight infants: is volume or pressure a better target variable? *J Pediatr* 2006; 3: 308–13.
230. Sinn JK, Ward MC, Henderson-Smart DJ. Developmental outcome of preterm infants after surfactant therapy: systematic review of randomized controlled trials. *J Paediatr Child Health* 2002; 6: 597–600.
231. Skippen P, Kissoon N. Ultrasound guidance for central vascular access in the pediatric emergency department. *Pediatr Emerg Care* 2007; 3: 203–7.
232. Sreenan C, Lemke RP, Hudson-Mason A, Osio-vich H. High-flow nasal cannulae in the management of apnea of prematurity: a comparison with conventional nasal continuous positive airway pressure. *Pediatrics* 2001; 5: 1081–3.
233. Stankiewicz P, Sen P, Bhatt SS et al. Genomic and genic deletions of the FOX gene cluster on 16q24.1 and inactivating mutations of FOXF1 cause alveolar capillary dysplasia and other malformations. *Am J Hum Genet* 2009; 6: 780–91.
234. Stenson B, Brocklehurst P, Tarnow-Mordi W. Increased 36-week survival with high oxygen saturation target in extremely preterm infants. *N Engl J Med* 2011; 17: 1680–2.
235. Stenson BJ, Boyle DW, Szylid EG. Initial ventilation strategies during newborn resuscitation. *Clin Perinatol* 2006; 1: 65–82, vi–vii.
236. Stevens TP, Blennow M, Soll RF. Early surfactant administration with brief ventilation vs selective surfactant and continued mechanical ventilation for preterm infants with or at risk for respiratory distress syndrome. *Cochrane Database Syst Rev* 2004; 3: CD003063.
237. Stevens TP, Harrington EW, Blennow M, Soll RF. Early surfactant administration with brief ventilation vs. selective surfactant and continued mechanical ventilation for preterm infants with or at risk for respiratory distress syndrome. *Cochrane Database Syst Rev* 2007; 4: CD003063.
238. Strohm B, Hobson A, Brocklehurst P et al. Subcutaneous fat necrosis after moderate therapeutic

- tic hypothermia in neonates. *Pediatrics* 2011; 2: e450–2.
239. Sud S, Sud M, Friedrich JO et al. High frequency oscillation in patients with acute lung injury and acute respiratory distress syndrome (ARDS): systematic review and meta-analysis. *BMJ* 2010; 340: c2327.
 240. Swarnam K, Soraisham AS, Sivanandan S. Advances in the management of meconium aspiration syndrome. *Int J Pediatr* 2012; 2012: 359571.
 241. Sweet DG, Carnielli V, Greisen G et al. (European Association of Perinatal Medicine). European consensus guidelines on the management of neonatal respiratory distress syndrome in preterm infants 2013 update. *Neonatology* 2013; 4: 353–68.
 242. The Neonatal Inhaled Nitric Oxide Study Group. Inhaled nitric oxide in term and near-term infants: neurodevelopmental follow-up of the neonatal inhaled nitric oxide study group (NINOS). *J Pediatr* 2000; 5: 611–7.
 243. Thompson CM, Puterman AS, Linley LL et al. The value of a scoring system for hypoxic ischaemic encephalopathy in predicting neurodevelopmental outcome. *Acta Paediatr* 1997; 7: 757–61.
 244. Thoresen M. Supportive care during neuroprotective hypothermia in the term newborn: adverse effects and their prevention. *Clin Perinatol* 2008; 4: 749–63, vii.
 245. Tin W, Milligan DW, Pennefather P, Hey E. Pulse oximetry, severe retinopathy, and outcome at one year in babies of less than 28 weeks gestation. *Arch Dis Child Fetal Neonatal Ed* 2001; 2: F106–10.
 246. Tin W, Walker S, Lacamp C. Oxygen monitoring in preterm babies: too high, too low? *Paediatr Respir Rev* 2003; 1: 9–14.
 247. Tobin R, Wetzel R. Cardiovascular Physiology and Shock. In: Nichols DG CD, Greeley WJ, Lappe DG, Underleider RM, Wetzel RC (Hrsg). *Cardiac intensive care in infants and children*. St. Louis: Mosby 1995; 17–74.
 248. Vain NE, Szyld EG, Prudent LM et al. Oropharyngeal and nasopharyngeal suctioning of meconium-stained neonates before delivery of their shoulders: multicentre, randomised controlled trial. *Lancet* 2004; 364: 597–602.
 249. Valls-i-Soler A, Lopez-Heredia J, Fernandez-Ruanova MB, Gastiasoro E. A simplified surfactant dosing procedure in respiratory distress syndrome: the »side-hole« randomized study. *Spanish Surfactant Collaborative Group. Acta Paediatr* 1997; 7: 747–51.
 250. Valmari P. Should pulse oximetry be used to screen for congenital heart disease? *Arch Dis Child Fetal Neonatal Ed* 2007; 3: F219–24.
 251. Van Marter LJ, Allred EN, Pagano M et al. Do clinical markers of barotrauma and oxygen toxicity explain interhospital variation in rates of chronic lung disease? The Neonatology Committee for the Developmental Network. *Pediatrics* 2000; 6: 1194–201.
 252. van Straaten HL, Rademaker CM, de Vries LS. Comparison of the effect of midazolam or vecuronium on blood pressure and cerebral blood flow velocity in the premature newborn. *Dev Pharmacol Ther* 1992; 4: 191–5.
 253. Vanpee M, Walfridsson-Schultz U, Katz-Salomon M et al. Resuscitation and ventilation strategies for extremely preterm infants: a comparison study between two neonatal centers in Boston and Stockholm. *Acta Paediatr Scand* 2007; 1: 10–6; discussion 8–9.
 254. Vento M, Asensi M, Sastre J et al. Resuscitation with room air instead of 100% oxygen prevents oxidative stress in moderately asphyxiated term neonates. *Pediatrics* 2001; 4: 642–7.
 255. Vento M, Asensi M, Sastre J et al. Oxidative stress in asphyxiated term infants resuscitated with 100% oxygen. *J Pediatr* 2003; 3: 240–6.
 256. Vento M, Sastre J, Asensi MA, Vina J. Room-air resuscitation causes less damage to heart and kidney than 100% oxygen. *Am J Respir Crit Care Med* 2005; 11: 1393–8.
 257. Verder H, Robertson B, Greisen G et al. Surfactant therapy and nasal continuous positive airway pressure for newborns with respiratory distress syndrome. Danish-Swedish Multicenter Study Group. *N Engl J Med* 1994; 16: 1051–5.
 258. Vergheze ST, McGill WA, Patel RI et al. Ultrasound-guided internal jugular venous cannulation in infants: a prospective comparison with the traditional palpation method. *Anesthesiology* 1999; 1: 71–7.
 259. Vohra S, Roberts RS, Zhang B et al. Heat loss prevention (HeLP) in the delivery room: a randomized controlled trial of polyethylene occlusive skin wrapping in very preterm infants. *J Pediatr* 2004; 145: 750–3.
 260. Ward M, Sinn J. Steroid therapy for meconium aspiration syndrome in newborn infants. *Cochrane Database Syst Rev* 2003; 4: CD003485.

261. Watterberg KL, Gerdes JS, Cole CH et al. Prophylaxis of early adrenal insufficiency to prevent bronchopulmonary dysplasia: a multicenter trial. *Pediatrics* 2004; 6: 1649–57.
 262. Watterberg KL, Shaffer ML, Mishefske MJ et al. Growth and neurodevelopmental outcomes after early low-dose hydrocortisone treatment in extremely low birth weight infants. *Pediatrics* 2007; 1: 40–8.
 263. Wilkinson D, Andersen C, O'Donnell CP, De Paoli AG. High flow nasal cannula for respiratory support in preterm infants. *Cochrane Database Syst Rev* 2011; 5: CD006405.
 264. Wiswell TE, Gannon CM, Jacob J et al. Delivery room management of the apparently vigorous meconium-stained neonate: results of the multicenter international collaborative trial. *Pediatrics* 2000; 1: 1–7.
 265. Wiswell TE, Knight GR, Finer NN et al. (Cochrane CG). A multicenter, randomized, controlled trial comparing Surfactin (Lucinactant) lavage with standard care for treatment of meconium aspiration syndrome. *Pediatrics* 2002; 6: 1081–7.
 266. Wyatt JS, Gluckman PD, Liu PY et al. Determinants of outcomes after head cooling for neonatal encephalopathy. *Pediatrics* 2007; 5: 912–21.
 267. Wyllie J, Perlman JM, Kattwinkel J et al. Part 11: Neonatal resuscitation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2010; 81 Suppl 1: e260–87. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
 268. Zhou WH, Cheng GQ, Shao XM et al. Selective head cooling with mild systemic hypothermia after neonatal hypoxic-ischemic encephalopathy: a multicenter randomized controlled trial in China. *J Pediatr* 2010; 3: 367–72, 72e1–3.
 269. Zola EM, Gunkel JH, Chan RK et al. Comparison of three dosing procedures for administration of bovine surfactant to neonates with respiratory distress syndrome. *J Pediatr* 1993; 3: 453–9.
- Teil III Typische Szenarien im Neugeborenenalter**
1. 2005 American Heart Association (AHA) guidelines for cardiopulmonary resuscitation (CPR) and emergency cardiovascular care (ECC) of pediatric and neonatal patients: neonatal resuscitation guidelines. *Pediatrics* 2006; 5: e1029–38.
 2. ACOG Practice Bulletin. Episiotomy. Clinical Management Guidelines for Obstetrician-Gynecologists. *Obstet Gynecol* 2006; 4: 957–62.
 3. AAP. The changing concept of sudden infant death syndrome: diagnostic coding shifts, controversies regarding the sleeping environment, and new variables to consider in reducing risk. *Pediatrics* 2005; 5: 1245–55.
 4. Abman SH (co-chair), Hansmann G (co-chair), Archer SL (co-chair) et al. AHA/ATS Joint Guidelines For Pediatric Pulmonary Hypertension. *Circulation* 2014 in press. Siehe hierzu auch special issue, expert consensus statement, European Pediatric Pulmonary Vascular Disease Network Heart (multipaper) 2015/2016 in press.
 5. ACOG Committee Opinion. Inappropriate use of the terms fetal distress and birth asphyxia. *Obstet Gynecol* 2005; 6: 1469–70.
 6. ACOG Committee Opinion. The Apgar score. *Obstet Gynecol* 2006; 5: 1209–12.
 7. Acosta-Rojas R, Becker J, Munoz-Abellana B et al. Twin chorionicity and the risk of adverse perinatal outcome. *Int J Gynaecol Obstet* 2007; 2: 98–102.
 8. Actelion. Persistent Pulmonary Hypertension of the Newborn (FUTURE 4). <https://clinicaltrials.gov/ct2/show/NCT01389856?term=01389856&rank=1>.
 9. Adamkin DH. Postnatal glucose homeostasis in late-preterm and term infants. *Pediatrics* 2011; 3: 575–9.
 10. Adzick NS. Management of fetal lung lesions. *Clin Perinatol* 2003; 3: 481–92.
 11. Adzick NS, Thom EA, Spong CY et al. A randomized trial of prenatal versus postnatal repair of myelomeningocele. *N Engl J Med* 2011; 11: 993–1004.
 12. Agthe AG, Kim GR, Mathias KB et al. Clonidine as an adjunct therapy to opioids for neonatal abstinence syndrome: a randomized, controlled trial. *Pediatrics* 2009; 5: e849–56.

13. Al-Salem AH. Congenital pyloric atresia and associated anomalies. *Pediatr Surg Int* 2007; 6: 559–63.
14. Albers LL, Anderson D, Cragin L et al. Factors related to perineal trauma in childbirth. *J Nurse Midwifery* 1996; 4: 269–76.
15. Alfaleh K, Anabrees J, Bassler D, Al-Kharfi T. Probiotics for prevention of necrotizing enterocolitis in preterm infants. *Cochrane Database Syst Rev* 2011; 3: CD005496.
16. Alkalay AL, Sarnat HB, Flores-Sarnat L et al. Population meta-analysis of low plasma glucose thresholds in full-term normal newborns. *Am J Perinatol* 2006; 2: 115–9.
17. Allen HD, Driscoll DJ, Shaddy RE, Feltes TF. Moss and Adams' heart disease in infants, children, and adolescents: including the fetus and young adult. Philadelphia: Lippincott Williams & Wilkins 2008.
18. Almubarak S, Wong PK. Long-term clinical outcome of neonatal EEG findings. *J Clin Neurophysiol* 2011; 2: 185–9.
19. Alsoufi B, Karamlou T, McCrindle BW, Caldarone CA. Management options in neonates and infants with critical left ventricular outflow tract obstruction. *Eur J Cardiothorac Surg* 2007; 6: 1013–21.
20. Alsoufi B, Bennetts J, Verma S, Caldarone CA. New developments in the treatment of hypoplastic left heart syndrome. *Pediatrics* 2007; 1: 109–17.
21. American Academy of Pediatrics. Group B streptococcal infections. In: *Red Book: report on infectious diseases*. Elk Grove Village, IL: American Academy of Pediatrics 2000.
22. Amiel J, Sproat-Emison E, Garcia-Barcelo M et al. Hirschsprung disease, associated syndromes and genetics: a review. *J Med Genet* 2008; 1: 1–14.
23. Anand RJ, Leaphart CL, Mollen KP, Hackam DJ. The role of the intestinal barrier in the pathogenesis of necrotizing enterocolitis. *Shock* 2007; 2: 124–33.
24. Andersson O, Hellstrom-Westas L, Andersson D, Domellof M. Effect of delayed versus early umbilical cord clamping on neonatal outcomes and iron status at 4 months: a randomised controlled trial. *BMJ* 2011; 343: d7157.
25. Armentrout DC, Huseby V. Polycythemia in the newborn. *MCN Am J Matern Child Nurs* 2003; 4: 234–9.
26. Askie LM, Ballard RA, Cutter GR et al. Meta-analysis of Preterm Patients on Inhaled Nitric Oxide C. Inhaled nitric oxide in preterm infants: an individual-patient data meta-analysis of randomized trials. *Pediatrics* 2011; 4: 729–39.
27. Azzopardi DV, Strohm B, Edwards AD et al. Moderate hypothermia to treat perinatal asphyxial encephalopathy. *N Engl J Med* 2009; 14: 1349–58.
28. Bagolan P, Nahom A, Giorlandino C et al. Cystic adenomatoid malformation of the lung: clinical evolution and management. *Eur J Pediatr* 1999; 11: 879–82.
29. Bahm J O-PC, Disselhorst-Klug C, Sellhaus B, Weis J. Obstetric brachial plexus palsy-treatment strategy, long-term results and prognosis. *Dtsch Arztebl Int* 2009; 106: 83–90.
30. Baird R, Eeson G, Safavi A et al. Canadian Pediatric Surgery N. Institutional practice and outcome variation in the management of congenital diaphragmatic hernia and gastroschisis in Canada: a report from the Canadian Pediatric Surgery Network. *J Pediatr Surg* 2011; 5: 801–7.
31. Baker PA, Brunette KE, Byrnes CA, Thompson JM. A prospective randomized trial comparing supraglottic airways for flexible bronchoscopy in children. *Paediatr Anaesth* 2010; 9: 831–8.
32. Ballabh P, Xu H, Hu F et al. Angiogenic inhibition reduces germinal matrix hemorrhage. *Nat Med* 2007; 4: 477–85.
33. Baquero H, Soliz A, Neira F et al. Oral sildenafil in infants with persistent pulmonary hypertension of the newborn: a pilot randomized blinded study. *Pediatrics* 2006; 4: 1077–83.
34. Barkovich AJ, Baranski K, Vigneron D et al. Proton MR spectroscopy for the evaluation of brain injury in asphyxiated, term neonates. *AJNR Am J Neuroradiol* 1999; 8: 1399–405.
35. Barrington KJ, Finer NN. Inhaled nitric oxide for preterm infants: a systematic review. *Pediatrics* 2007; 5: 1088–99.
36. Barrington KJ, Finer NN. Inhaled nitric oxide for respiratory failure in preterm infants. *Cochrane Database Syst Rev* 2007; 3: CD000509.
37. Barrington KJ, Finer N. Inhaled nitric oxide for respiratory failure in preterm infants. *Cochrane Database Syst Rev* 2010; 12: CD000509.
38. Bastian H, Keirse MJ, Lancaster PA. Perinatal death associated with planned home birth in

- Australia: population based study. *BMJ* 1998; 7155: 384–8.
39. Batton B, Batton D, Riggs T. Blood pressure during the first 7 days in premature infants born at postmenstrual age 23 to 25 weeks. *Am J Perinatol* 2007; 2: 107–15.
 40. Beardsall K. Measurement of glucose levels in the newborn. *Early human development* 2010; 5: 263–7.
 41. Bedwell J, Shah RK, Bauman N, Zalzal GH, Preciado DA. Balloon dilation for management of choanal atresia and stenosis. *Int J Pediatr Otorhinolaryngol* 2011; 12: 1515–8.
 42. Beghetti M, La Scala G, Belli D et al. Etiology and management of pediatric chylothorax. *J Pediatr* 2000; 5: 653–8.
 43. Bellini C, Hennekam RC, Fulcheri E et al. Etiology of nonimmune hydrops fetalis: a systematic review. *Am J Med Genet A* 2009; 5: 844–51.
 44. Ben-Ari YaHG. Effects of seizures on developmental processes in the immature brain. *Lancet Neurol* 2006; 12: 1055–63.
 45. Benitz WE, Han MY, Madan A, Ramachandra P. Serial serum C-reactive protein levels in the diagnosis of neonatal infection. *Pediatrics* 1998; 4: E41.
 46. Berger TM, Das-Kundu S, Pfister RE et al. Prevention and therapy of hypoglycemia in infants with a gestational age above 34 0/7 weeks in maternity wards. Recommendations by the Swiss Society of Neonatology 2007.
 47. Berman L, Moss RL. Necrotizing enterocolitis: an update. *Semin Fetal Neonatal Med* 2011; 3: 145–50.
 48. Berner R. Group B streptococci during pregnancy and infancy. *Curr Opin Infect Dis* 2002; 3: 307–13.
 49. Bhuta T, Henderson-Smart DJ. Rescue high frequency oscillatory ventilation versus conventional ventilation for pulmonary dysfunction in preterm infants. *Cochrane Database Syst Rev* 2000; 2: CD000438.
 50. Bianca S, Barbagallo MA, Ingegnesi C, Ettore G. Isolated hypertrophic pyloric stenosis and perinatal factors. *Genet Couns* 2003; 1: 101–3.
 51. Birdi K, Prasad AN, Prasad C et al. The floppy infant: retrospective analysis of clinical experience (1990–2000) in a tertiary care facility. *J Child Neurol* 2005; 10: 803–8.
 52. Bittigau P, Sifringer M, Genz K et al. Antiepileptic drugs and apoptotic neurodegeneration in the developing brain. *Proc Natl Acad Sci U S A* 2002; 23: 15089–94.
 53. Blakely ML, Tyson JE, Lally KP et al. Laparotomy versus peritoneal drainage for necrotizing enterocolitis or isolated intestinal perforation in extremely low birth weight infants: outcomes through 18 months adjusted age. *Pediatrics* 2006; 4: e680–7.
 54. Blakely ML, Gupta H, Lally KP. Surgical management of necrotizing enterocolitis and isolated intestinal perforation in premature neonates. *Semin Perinatol* 2008; 2: 122–6.
 55. Boloker J, Bateman DA, Wung JT, Stolar CJ. Congenital diaphragmatic hernia in 120 infants treated consecutively with permissive hypercapnea/spontaneous respiration/elective repair. *J Pediatr Surg* 2002; 3: 357–66.
 56. Boluyt N, van Kempen A, Offringa M. Neurodevelopment after neonatal hypoglycemia: a systematic review and design of an optimal future study. *Pediatrics* 2006; 6: 2231–43.
 57. Booth D, Evans DJ. Anticonvulsants for neonates with seizures. *Cochrane Database Syst Rev* 2004; 4: CD004218.
 58. Bose CL, Laughon M. Treatment to prevent patency of the ductus arteriosus: beneficial or harmful? *J Pediatr* 2006; 6: 713–4.
 59. Bose CL, Laughon MM. Patent ductus arteriosus: lack of evidence for common treatments. *Arch Dis Child Fetal Neonatal Ed* 2007; 6: F498–502.
 60. Bouissou A, Rakza T, Klosowski S et al. Hypotension in preterm infants with significant patent ductus arteriosus: effects of dopamine. *J Pediatr* 2008; 6: 790–4.
 61. Brown JW, Ruzmetov M, Okada Y et al. Outcomes in patients with interrupted aortic arch and associated anomalies: a 20-year experience. *Eur J Cardiothorac Surg* 2006; 5: 666–74.
 62. Bühner C, Bahr S, Siebert J et al. Use of 2% 2-phenoxyethanol and 0.1% octenidine as antiseptic in premature newborn infants of 23–26 weeks gestation. *J Hosp Infect* 2002; 4: 305–7.
 63. Bullock R, Martin WL, Coomarasamy A, Kilby MD. Prediction of fetal anemia in pregnancies with red-cell alloimmunization: comparison of middle cerebral artery peak systolic velocity and amniotic fluid OD450. *Ultrasound Obstet Gynecol* 2005; 4: 331–4.
 64. Burge D, Drewett M. Meconium plug obstruction. *Pediatr Surg Int* 2004; 2: 108–10.

65. Burns CM, Rutherford MA, Boardman JP, Cowan FM. Patterns of cerebral injury and neurodevelopmental outcomes after symptomatic neonatal hypoglycemia. *Pediatrics* 2008; 1: 65–74.
66. Burri S, Hug MI, Bauersfeld U. Efficacy and safety of intravenous amiodarone for incessant tachycardias in infants. *Eur J Pediatr* 2003; 12: 880–4.
67. Buyon JP, Hiebert R, Copel J et al. Autoimmune-associated congenital heart block: demographics, mortality, morbidity and recurrence rates obtained from a national neonatal lupus registry. *J Am Coll Cardiol* 1998; 7: 1658–66.
68. Buyon JP, Rupel A, Clancy RM. Neonatal lupus syndromes. *Lupus* 2004; 9: 705–12.
69. Calandre EP, Dominguez-Granados R, Gomez-Rubio M, Molina-Font JA. Cognitive effects of long-term treatment with phenobarbital and valproic acid in school children. *Acta Neurol Scand* 1990; 6: 504–6.
70. Camfield CS, Chaplin S, Doyle AB et al. Side effects of phenobarbital in toddlers; behavioral and cognitive aspects. *J Pediatr* 1979; 3: 361–5.
71. Carter BS, Haverkamp AD, Merenstein GB. The definition of acute perinatal asphyxia. *Clin Perinatol* 1993; 2: 287–304.
72. Case AP, Mitchell LE. Prevalence and patterns of choanal atresia and choanal stenosis among pregnancies in Texas, 1999–2004. *Am J Med Genet A* 2011; 4: 786–91.
73. Cerro MJ, Abman S, Diaz G et al. A consensus approach to the classification of pediatric pulmonary hypertensive vascular disease: Report from the PVRI Pediatric Taskforce, Panama 2011. *Pulm Circ* 2011; 2: 286–98.
74. Chalouhi GE, Stirnemann JJ, Salomon LJ et al. Specific complications of monochorionic twin pregnancies: twin-twin transfusion syndrome and twin reversed arterial perfusion sequence. *Semin Fetal Neonatal Med* 2010; 6: 349–56.
75. Chalouhi GE, Essaoui M, Stirnemann J et al. Laser therapy for twin-to-twin transfusion syndrome (TTTS). *Prenat Diagn* 2011; 7: 637–46.
76. Chambers CD, Hernandez-Diaz S, Van Marter LJ et al. Selective serotonin-reuptake inhibitors and risk of persistent pulmonary hypertension of the newborn. *N Engl J Med* 2006; 6: 579–87.
77. Chamnanvanakij S, Rollins N, Perlman JM. Subdural hematoma in term infants. *Pediatr Neurol* 2002; 4: 301–4.
78. Chang DC, Salazar-Osuna JH, Choo SS et al. Benchmarking the quality of care of infants with low-risk gastroschisis using a novel risk stratification index. *Surgery* 2010; 6: 766–71.
79. Charney EB, Melchionni JB, Antonucci DL. Ventriculitis in newborns with myelomeningocele. *Am J Dis Child* 1991; 3: 287–90.
80. Cheung YF, Leung MP, Lee JW et al. Evolving management for critical pulmonary stenosis in neonates and young infants. *Cardiol Young* 2000; 3: 186–92.
81. Choi HJ, Hahn S, Lee J et al. Surfactant lavage therapy for meconium aspiration syndrome: a systematic review and meta-analysis. *Neonatology* 2012; 3: 183–91.
82. Christison-Lagay ER, Kelleher CM, Langer JC. Neonatal abdominal wall defects. *Semin Fetal Neonatal Med* 2011; 3: 164–72.
83. Clancy RR. Summary proceedings from the neurology group on neonatal seizures. *Pediatrics* 2006; 3 Pt 2: S23–7.
84. Clyman RI. Ibuprofen and patent ductus arteriosus. *N Engl J Med* 2000; 10: 728–30.
85. Clyman RI, Saha S, Jobe A, Oh W. Indomethacin prophylaxis for preterm infants: the impact of 2 multicentered randomized controlled trials on clinical practice. *J Pediatr* 2007; 1: 46–50e2.
86. Cohen LS, Altshuler LL, Harlow BL, Nonacs R, Newport DJ, Viguera AC, Suri R, Burt VK, Hendrick V, Reminick AM, Loughhead A, Vitonis AF, Stowe ZN. Relapse of major depression during pregnancy in women who maintain or discontinue antidepressant treatment. *JAMA* 2006; 5: 499–507.
87. Cohen-Overbeek TE, Tong WH, Hatzmann TR et al. Omphalocele: comparison of outcome following prenatal or postnatal diagnosis. *Ultrasound Obstet Gynecol* 2010; 6: 687–92.
88. Committee on Infectious D. From the American Academy of Pediatrics: Policy statements – Modified recommendations for use of palivizumab for prevention of respiratory syncytial virus infections. *Pediatrics* 2009; 6: 1694–701.
89. Connor JA, Thiagarajan R. Hypoplastic left heart syndrome. *Orphanet J Rare Dis* 2007; 2: 23.
90. Cornblath M, Hawdon JM, Williams AF et al. Controversies regarding definition of neonatal hypoglycemia: suggested operational thresholds. *Pediatrics* 2000; 5: 1141–5.

91. Corrales CE, Koltai PJ. Choanal atresia: current concepts and controversies. *Curr Opin Otolaryngol Head Neck Surg* 2009; 17: 466–70.
92. Cotten CM. Air leak syndromes. In: Spitzer A (Hrsg). *Intensive Care of the Fetus and Neonate*. Philadelphia: Elsevier Mosby 2005.
93. Cowles RA, Berdon WE, Holt PD et al. Neonatal intestinal obstruction simulating meconium ileus in infants with long-segment intestinal aganglionosis: radiographic findings that prompt the need for rectal biopsy. *Pediatr Radiol* 2006; 2: 133–7.
94. Cua CL, Blankenship A, North AL et al. Increased incidence of idiopathic persistent pulmonary hypertension in down syndrome neonates. *Pediatr Cardiol* 2007; 4: 250–4.
95. Curran PF, Jelin EB, Rand L et al. Prenatal steroids for microcystic congenital cystic adenomatoid malformations. *J Pediatr Surg* 2010; 1: 145–50.
96. Czernik C, Proquitté H, Metze B, Bühner C. Hydrops fetalis – has there been a change in diagnostic spectrum and mortality? *J Matern Fetal Neonatal Med* 2011; 2: 258–63.
97. Danzer E, Hedrick HL. Neurodevelopmental and neurofunctional outcomes in children with congenital diaphragmatic hernia. *Early Hum Dev* 2011; 9: 625–32.
98. Dargaville PA, Copnell B, Mills JF et al. Randomized controlled trial of lung lavage with dilute surfactant for meconium aspiration syndrome. *J Pediatr* 2011; 3: 383–9e2.
99. Davenport M, Warne SA, Cacciaguerra S et al. Current outcome of antenally diagnosed cystic lung disease. *J Pediatr Surg* 2004; 4: 549–56.
100. Dawson JA, Kamlin CO, Wong C et al. Changes in heart rate in the first minutes after birth. *Arch Dis Child Fetal Neonatal Ed* 2010; 3: F177–81.
101. Dawson JA, Kamlin CO, Vento M et al. Defining the reference range for oxygen saturation for infants after birth. *Pediatrics* 2010; 6: e1340–7.
102. De Jonge A, van der Goes BY, Ravelli AC et al. Perinatal mortality and morbidity in a nationwide cohort of 529,688 low-risk planned home and hospital births. *BJOG* 2009; 9: 1177–84.
103. De la Hunt MN. The acute abdomen in the newborn. *Semin Fetal Neonatal Med* 2006; 3: 191–7.
104. De Vries LS, Toet MC. Amplitude integrated electroencephalography in the full-term newborn. *Clin Perinatol* 2006; 3: 619–32.
105. De Wals P, Tairou F, Van Allen MI et al. Reduction in neural-tube defects after folic acid fortification in Canada. *N Engl J Med* 2007; 2: 135–42.
106. De Wolf D, Vanderbruggen K, Verbist A et al. Percutaneous interventions for congenital aortic stenosis. *Acta Cardiol* 2006; 2: 204–5.
107. DeBord S, Cherry C, Hickey C. The arterial switch procedure for transposition of the great arteries. *AORN J* 2007; 2: 211–30.
108. Delaney C, Cornfield DN. Risk factors for persistent pulmonary hypertension of the newborn. *Pulm Circ* 2012; 1: 15–20.
109. DeMello DE. Pulmonary pathology. *Semin Neonatol* 2004; 4: 311–29.
110. Dempsey EM, Barrington K. Crystalloid or colloid for partial exchange transfusion in neonatal polycythemia: a systematic review and meta-analysis. *Acta Paediatr Scand* 2005; 11: 1650–5.
111. Derby CD, Pizarro C. Routine primary repair of tetralogy of Fallot in the neonate. *Expert Rev Cardiovasc Ther* 2005; 3: 857–63.
112. Deshpande G, Rao S, Patole S, Bulsara M. Updated meta-analysis of probiotics for preventing necrotizing enterocolitis in preterm neonates. *Pediatrics* 2010; 5: 921–30.
113. Deshpande GC, Rao SC, Keil AD, Patole SK. Evidence-based guidelines for use of probiotics in preterm neonates. *BMC Medicine* 2011; 9: 92.
114. Deutsche Gesellschaft für Kinderchirurgie. Kurzstreckige Ösophagusatresie (AWMF-Register Nr. 006/045; www.awmf.org).
115. Deutsche Gesellschaft für Schlafmedizin. Plötzlicher Säuglingstod. (AWMF-Register Nr. 063/002; www.awmf.org).
116. Diaz J, Schain RJ. Phenobarbital: effects of long-term administration on behavior and brain of artificially reared rats. *Science* 1978; 199: 90–1.
117. Dimmitt RA, Moss RL. Meconium diseases in infants with very low birth weight. *Semin Pediatr Surg* 2000; 9: 79–83.
118. Dixon G, Badawi N, Kurinczuk JJ et al. Early developmental outcomes after newborn encephalopathy. *Pediatrics* 2002; 1: 26–33.
119. Duran R, Aladag N, Vatansever U et al. The impact of Neonatal Resuscitation Program courses on mortality and morbidity of newborn infants with perinatal asphyxia. *Brain Dev* 2008; 1: 43–6.
120. Duvanel CB, Fawer CL, Cotting J et al. Long-term effects of neonatal hypoglycemia on brain growth and psychomotor development in small-

- for-gestational-age preterm infants. *J Pediatr* 1999; 4: 492–8.
121. Dzhala VI, Talos DM, Sdrulla DA et al. NKCC1 transporter facilitates seizures in the developing brain. *Nat Med* 2005; 11: 1205–13.
 122. Echtler K, Stark K, Lorenz M et al. Platelets contribute to postnatal occlusion of the ductus arteriosus. *Nat Med* 2010; 1: 75–82.
 123. Edwards AD, Azzopardi DV. Therapeutic hypothermia following perinatal asphyxia. *Arch Dis Child Fetal Neonatal Ed* 2006; 2: F127–31.
 124. Edwards JE, Burchell HB. Congenital tricuspid atresia; a classification. *Med Clin North Am* 1949; 33: 1177–96.
 125. Eicher C, Seitz G, Bevot A et al. Surgical management of extremely low birth weight infants with neonatal bowel perforation: a single-center experience and a review of the literature. *Neonatology* 2012; 4: 285–92.
 126. Eicher DJ, Wagner CL, Katikaneni LP et al. Moderate hypothermia in neonatal encephalopathy: safety outcomes. *Pediatr Neurol* 2005; 1: 18–24.
 127. Ein SH, Palder SB, Filler RM. Babies with esophageal and duodenal atresia: a 30-year review of a multifaceted problem. *J Pediatr Surg* 2006; 3: 530–2.
 128. El Shahed A, Dargaville P, Ohlsson A, Soll R. Surfactant for meconium aspiration syndrome in full term/near term infants. *Cochrane Database Syst Rev* 2007; 3: CD002054.
 129. Epelman M. The whirlpool sign. *Radiology* 2006; 3: 910–1.
 130. Erler T, Beyer U, Hoch B et al. Heimüberwachung (»home monitoring«) von Kindern und Jugendlichen: Vorschläge für die praktische Anwendung. *Somnologie* 2009; 3: 182–8.
 131. Eschenbach DA. Prevention of neonatal group B streptococcal infection. *N Engl J Med* 2002; 4: 280–1.
 132. Evans D, Levene M. Hypoxic-ischemic injury. Edinburgh: Churchill Livingstone 1999.
 133. Faingold R, Daneman A, Tomlinson G et al. Necrotizing enterocolitis: assessment of bowel viability with color doppler US. *Radiology* 2005; 2: 587–94.
 134. Farrow KN, Groh BS, Schumacker PT et al. Hyperoxia increases phosphodiesterase 5 expression and activity in ovine fetal pulmonary artery smooth muscle cells. *Circ Res* 2008; 2: 226–33.
 135. Fasnacht M, Pfammatter JP, Ghisla R et al. 40th Annual Meeting of the AEPC 2005.
 136. Ferencz C, Rubin JD, McCarter RJ et al. Congenital heart disease: prevalence at livebirth. The Baltimore-Washington Infant Study. *Am J Epidemiol* 1985; 1: 31–6.
 137. Filippi L, Pezzati M, Poggi C et al. Dopamine versus dobutamine in very low birthweight infants: endocrine effects. *Arch Dis Child Fetal Neonatal Ed* 2007; 5: F367–71.
 138. Findlay RD, Taeusch HW, Walther FJ. Surfactant replacement therapy for meconium aspiration syndrome. *Pediatrics* 1996; 1: 48–52.
 139. Finer NN, Barrington KJ. Nitric oxide for respiratory failure in infants born at or near term. *Cochrane Database Syst Rev* 2006; 4: CD000399.
 140. Finer NN, Carlo WA, Walsh MC et al. Early CPAP versus surfactant in extremely preterm infants. *N Engl J Med* 2010; 21: 1970–9.
 141. Fitzgibbons SC, Ching Y, Yu D et al. Mortality of necrotizing enterocolitis expressed by birth weight categories. *J Pediatr Surg* 2009; 6: 1072–6.
 142. Fleming P, Blair PS. Sudden Infant Death Syndrome and parental smoking. *Early Hum Dev* 2007; 11: 721–5.
 143. Fligor BJ, Neault MW, Mullen CH et al. Factors associated with sensorineural hearing loss among survivors of extracorporeal membrane oxygenation therapy. *Pediatrics* 2005; 6: 1519–28.
 144. Foker JE, Linden BC, Boyle EM Jr, Marquardt C. Development of a true primary repair for the full spectrum of esophageal atresia. *Ann Surg* 1997; 4: 533–43.
 145. Fowlie PW, Davis PG, McGuire W. Prophylactic intravenous indomethacin for preventing mortality and morbidity in preterm infants. *Cochrane Database Syst Rev* 2010; 7: CD000174.
 146. Frisk V, Jakobson LS, Unger S et al. Long-term neurodevelopmental outcomes of congenital diaphragmatic hernia survivors not treated with extracorporeal membrane oxygenation. *J Pediatr Surg* 2011; 7: 1309–18.
 147. Fukushima K, Morokuma S, Fujita Y et al. Short-term and long-term outcomes of 214 cases of non-immune hydrops fetalis. *Early Hum Dev* 2011; 8: 571–5.
 148. Fyler DC. Report of the New England Regional Infant Cardiac Program. *Pediatrics* 1980; 65 (Suppl): 375–461.
 149. Gaede C. Congenital chylothorax: a case study. *Neonatal Netw* 2006; 5: 371–81.

150. Gandy GM, Adamsons K Jr, Cunningham N et al. Thermal Environment and Acid-Base Homeostasis in Human Infants during the First Few Hours of Life. *J Clin Invest* 1964; 43: 751–8.
151. Gilbert D, Moellering R, Sande M. The Sanford guide to antimicrobial therapy. Hyde Park VT 2001.
152. Giunta RE, Enders A, Lukas B et al. Geburtstraumatische Armplexusparesen – Eine interdisziplinäre Herausforderung. *Monatsschr Kinderheilk* 2010; 158: 262–72.
153. Gluckman PD, Wyatt JS, Azzopardi D et al. Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicentre randomised trial. *Lancet* 2005; 365: 663–70.
154. Goldman LE, Boramanand NK, Acevedo V, Gallagher P, Nehgme R. Preterm infants with paroxysmal supraventricular tachycardia: presentation, response to therapy, and outcome. *J Interv Card Electrophysiol* 2001; 3: 293–7.
155. Gopel W, Kribs A, Ziegler A et al. Avoidance of mechanical ventilation by surfactant treatment of spontaneously breathing preterm infants (AMV): an open-label, randomised, controlled trial. *Lancet* 2011; 378: 1627–34.
156. Gordon PA. Congenital heart block: clinical features and therapeutic approaches. *Lupus* 2007; 8: 642–6.
157. Gottstein R, Cooke RW. Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn. *Arch Dis Child Fetal Neonatal Ed* 2003; 1: F6–10.
158. Goyal A, Manalang LR, Donnell SC, Lloyd DA. Primary peritoneal drainage in necrotising enterocolitis: an 18-year experience. *Pediatr Surg Int* 2006; 5: 449–52.
159. Greenough A, Sharma A. What is new in ventilation strategies for the neonate? *Eur J Pediatr* 2007; 10: 991–6.
160. Grifka RG. Cyanotic congenital heart disease with increased pulmonary blood flow. *Pediatr Clin North Am* 1999; 2: 405–25.
161. Grosch-Worner I, Schafer A, Obladen M et al. An effective and safe protocol involving zidovudine and caesarean section to reduce vertical transmission of HIV-1 infection. *AIDS* 2000; 18: 2903–11.
162. Group EM. Changing Child birth. HMSO 1993.
163. Gupta AK, Guglani B. Imaging of congenital anomalies of the gastrointestinal tract. *Indian J Pediatr* 2005; 5: 403–14.
164. Guthmann F, Bühner C. Routine probiotics in preterm infants? *Arch Dis Child Fetal Neonatal Ed* 2011; 4: F311–2.
165. Hagberg H, Wennerholm UB, Savman K. Sequelae of chorioamnionitis. *Curr Opin Infect Dis* 2002; 3: 301–6.
166. Hahn S, Choi HJ, Soll R, Dargaville PA. Lung lavage for meconium aspiration syndrome in newborn infants. *Cochrane Database Syst Rev* 2013; 4: CD003486.
167. Hall NJ, Curry J, Drake DP et al. Resection and primary anastomosis is a valid surgical option for infants with necrotizing enterocolitis who weigh less than 1000g. *Arch Surg* 2005; 12: 1149–51.
168. Hallberg B, Grossmann K, Bartocci M, Blenow M. The prognostic value of early aEEG in asphyxiated infants undergoing systemic hypothermia treatment. *Acta paediatr* 2010; 4: 531–6.
169. Hamrick SE, Hansmann G. Patent ductus arteriosus of the preterm infant. *Pediatrics* 2010; 5: 1020–30.
170. Hancock Friesen CL, Zurakowski D, Thiagarajan RR, Forbess JM, del Nido PJ, Mayer JE, Jonas RA. Total anomalous pulmonary venous connection: an analysis of current management strategies in a single institution. *Ann Thorac Surg* 2005; 2: 596–606.
171. Harvey D, Holt DE, Bedford H. Bacterial meningitis in the newborn: a prospective study of mortality and morbidity. *Semin Perinatol* 1999; 3: 218–25.
172. Hawdon JM. Investigation and management of impaired metabolic adaptation presenting as neonatal hypoglycaemia. *Paediatrics and Child Health* 2008; 4: 161–5.
173. Hemming V, Rankin J. Small intestinal atresia in a defined population: occurrence, prenatal diagnosis and survival. *Prenat Diagn* 2007; 27: 1205–11.
174. Herrera C, Holberton J, Davis P. Prolonged versus short course of indomethacin for the treatment of patent ductus arteriosus in preterm infants. *Cochrane Database Syst Rev* 2007; 2: CD003480.
175. Giagnorio R, Deindl P, Maiweg E et al. Persistierende pulmonale Hypertonie des Neugeborenen und chronisch progressive pulmonale Hypertonie im ersten Lebensjahr. *Kinder- und Jugendmedizin* 2015/2016 (im Druck).
176. Hoehn T, Hansmann G, Bühner C, Simbruner G, Gunn AJ, Yager J, Levene M, Hamrick SE, Shan-

- karan S, Thoresen M. Therapeutic hypothermia in neonates. Review of current clinical data, ILCOR recommendations and suggestions for implementation in neonatal intensive care units. *Resuscitation* 2008; 1: 7–12.
177. Hoepfer MM, Schwarze M, Ehlerding S, Adler-Schuermeier A, Spiekerkoetter E, Niedermeyer J, Hamm M, Fabel H. Long-term treatment of primary pulmonary hypertension with aerosolized iloprost, a prostacyclin analogue. *N Engl J Med* 2000; 25: 1866–70.
 178. Hofman FN, Bax NM, van der Zee DC, Kramer WL. Surgery for necrotising enterocolitis: primary anastomosis or enterostomy? *Pediatr Surg Int* 2004; 7: 481–3.
 179. Hofmeyr GJ, Lawrie TA. Amnioinfusion for potential or suspected umbilical cord compression in labour. *Cochrane Database Syst Rev* 2012; 1: CD000013.
 180. Holschneider A, Hutson J, Pena A et al. Preliminary report on the International Conference for the Development of Standards for the Treatment of Anorectal Malformations. *J Pediatr Surg* 2005; 10: 1521–6.
 181. Holt PD, Arkovitz MS, Berdon WE, Stolar CJ. Newborns with diaphragmatic hernia: initial chest radiography does not have a role in predicting clinical outcome. *Pediatr Radiol* 2004; 6: 462–4.
 182. Horvers M, Mooij CF, Antonius TA. Is octreotide treatment useful in patients with congenital chylothorax? *Neonatology* 2012; 3: 225–31.
 183. Hudak ML, Tan RC. Neonatal Drug Withdrawal. *Pediatrics* 2012; 2: e540–60.
 184. Hudgins RJ, Gilreath CL. Tethered spinal cord following repair of myelomeningocele. *Neurosurg Focus* 2004; 2: E7.
 185. Humpl T, Soderberg B, McCrindle BW et al. Percutaneous balloon valvotomy in pulmonary atresia with intact ventricular septum: impact on patient care. *Circulation* 2003; 7: 826–32.
 186. Humpl T, Reyes JT, Holtby H et al. Beneficial effect of oral sildenafil therapy on childhood pulmonary arterial hypertension: twelve-month clinical trial of a single-drug, open-label, pilot study. *Circulation* 2005; 24: 3274–80.
 187. Hung KL, Liao HT, Huang JS. Rational management of simple depressed skull fractures in infants. *J Neurosurg* 2005; 1 Suppl: 69–72.
 188. Hunt RW, Neil JJ, Coleman LT et al. Apparent diffusion coefficient in the posterior limb of the internal capsule predicts outcome after perinatal asphyxia. *Pediatrics* 2004; 4: 999–1003.
 189. Hüsemann D, Metze B, Walch E, Bühner C. Laboratory markers of perinatal acidosis are poor predictors of neurodevelopmental impairment in very low birth weight infants. *Early Hum Dev* 2011; 10: 677–81.
 190. Ichinose F, Erana-Garcia J, Hromi J et al. Nebulized sildenafil is a selective pulmonary vasodilator in lambs with acute pulmonary hypertension. *Crit Care Med* 2001; 5: 1000–5.
 191. Isemann B, Meinzen-Derr J, Akinbi H. Maternal and neonatal factors impacting response to methadone therapy in infants treated for neonatal abstinence syndrome. *J Perinatol* 2011; 1: 25–9.
 192. Ivy DD, Abman SH, Barst RJ et al. Pediatric pulmonary hypertension. *J Am Coll Cardiol* 2013; 25 Suppl: D117–26. Siehe hierzu auch special issue, expert consensus statement, European Pediatric Pulmonary Vascular Disease Network Heart (multipaper) 2015/2016 in press.
 193. Izmirly PM, Rivera TL, Buyon JP. Neonatal lupus syndromes. *Rheum Dis Clin North Am* 2007; 2: 267–85, vi.
 194. Jansson LM, Choo R, Velez ML et al. Methadone maintenance and breastfeeding in the neonatal period. *Pediatrics* 2008; 1: 106–14.
 195. Jensen FE. The role of glutamate receptor maturation in perinatal seizures and brain injury. *Int J Dev Neurosci* 2002; 20: 339–47.
 196. Jentink J, Loane MA, Dolk H et al. Valproic acid monotherapy in pregnancy and major congenital malformations. *N Engl J Med* 2010; 23: 2185–93.
 197. Johns Hopkins Hospital. Engorn B, Flerlage J. *The Harriet Lane Handbook: Mobile Medicine Series* (20th ed). Expert Consult. Philadelphia: Saunders by Elsevier 2015.
 198. Johnson BA, Ades A. Delivery room and early postnatal management of neonates who have prenatally diagnosed congenital heart disease. *Clin Perinatol* 2005; 4: 921–46, ix.
 199. Johnson WH Jr, Dunnigan A, Fehr P, Benson DW Jr. Association of atrial flutter with orthodromic reciprocating fetal tachycardia. *Am J Cardiol* 1987; 4: 374–5.
 200. Jones HE, Kaltenbach K, Heil SH et al. Neonatal abstinence syndrome after methadone or buprenorphine exposure. *N Engl J Med* 2010; 24: 2320–31.

201. Kaiser JR, Gauss CH, Pont MM, Williams DK. Hypercapnia during the first 3 days of life is associated with severe intraventricular hemorrhage in very low birth weight infants. *J Perinatol* 2006; 5: 279–85.
202. Kajimoto H, Hashimoto K, Bonnet SN et al. Oxygen activates the Rho/Rho-kinase pathway and induces RhoB and ROCK-1 expression in human and rabbit ductus arteriosus by increasing mitochondria-derived reactive oxygen species: a newly recognized mechanism for sustaining ductal constriction. *Circulation* 2007; 116: 1777–88.
203. Kallen BA, Otterblad Olausson P, Danielsson BR. Is erythromycin therapy teratogenic in humans? *Reprod Toxicol* 2005; 2: 209–14.
204. Kanegaye JT, Soliemanzadeh P, Bradley JS. Lumbar puncture in pediatric bacterial meningitis: defining the time interval for recovery of cerebrospinal fluid pathogens after parenteral antibiotic pretreatment. *Pediatrics* 2001; 107: 1169–74.
205. Kato I, Groswasser J, Franco P et al. Developmental characteristics of apnea in infants who succumb to sudden infant death syndrome. *Am J Respir Crit Care Med* 2001; 164: 1464–9.
206. Kato I, Franco P, Groswasser J et al. Incomplete arousal processes in infants who were victims of sudden death. *Am J Respir Crit Care Med* 2003; 168: 1298–303.
207. Kattner E, Maier RF, Gonser M et al. Betreuung von Neugeborenen diabetischer Mütter (AWMF-Register Nr. 024/006; www.awmf.org).
208. Kattwinkel J. Surfactant lavage for meconium aspiration: a word of caution. *Pediatrics* 2002; 110: 1167–8.
209. Kattwinkel J, Perlman JM, Aziz K et al. Part 15: neonatal resuscitation: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010; 122 Suppl 3: S909–19.
210. Kibar Z, Salem S, Bosoi CM et al. Contribution of VANGL2 mutations to isolated neural tube defects. *Clin Genet* 2011; 1: 76–82.
211. Kilani RA, Wetmore J. Neonatal subgaleal hematoma: presentation and outcome – radiological findings and factors associated with mortality. *Am J Perinatol* 2006; 1: 41–8.
212. Kimura K, Soper RT. Multistaged extrathoracic esophageal elongation for long gap esophageal atresia. *J Pediatr Surg* 1994; 4: 566–8.
213. Klebermass K, Olischar M, Waldhoer T, Fuiiko R, Pollak A, Weninger M. Amplitude-integrated EEG pattern predicts further outcome in preterm infants. *Pediatr Res* 2011; 1: 102–8.
214. Kleinman ME, Chameides L, Schexnayder SM et al. Part 14: pediatric advanced life support: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010; 122 Suppl 3: S876–908. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation* and *Resuscitation*.
215. Kliegman R. Fetal and neonatal medicine. In: Behrman R, Kliegman R (eds). *Nelson Essentials of Pediatrics*. Philadelphia: W. R. Saunders Company 2002; 179–249.
216. Kliegmann RM, Stanton B, Geme JS, Schor NF. *Nelson Notebook of Pediatrics*. 2-Volume set. 20th ed. Santa Barbara: Elsevier 2014.
217. Knight DB. The treatment of patent ductus arteriosus in preterm infants. A review and overview of randomized trials. *Semin Neonatol* 2001; 1: 63–73.
218. Kovalchin JP, Silverman NH. The impact of fetal echocardiography. *Pediatr Cardiol* 2004; 3: 299–306.
219. Kovesi T, Rubin S. Long-term complications of congenital esophageal atresia and/or tracheo-esophageal fistula. *Chest* 2004; 3: 915–25.
220. Krapp M, Ludwig A, Axt-Fliedner R, Kreiselmaier P. First trimester fetal echocardiography: which planes and defects can be displayed during the daily routine in a prenatal medicine unit? *Ultraschall Med* 2011; 4: 362–6.
221. Kribs A, Vierzig A, Hunseler C et al. Early surfactant in spontaneously breathing with nCPAP in ELBW infants – a single centre four year experience. *Acta Paediatr* 2008; 3: 293–8.
222. Kumaran N, Shankar KR, Lloyd DA, Losty PD. Trends in the management and outcome of jejuno-ileal atresia. *Eur J Pediatr Surg* 2002; 3: 163–7.
223. Lally KP, Lally PA, Van Meurs KP et al. Treatment evolution in high-risk congenital diaphragmatic hernia: ten years' experience with diaphragmatic agenesis. *Ann Surg* 2006; 4: 505–13.
224. Lansdale N, Hill R, Gull-Zamir S et al. Staged reduction of gastroschisis using preformed silos: practicalities and problems. *J Pediatr Surg* 2009; 44: 2126–9.

225. Lanska MJ, Lanska DJ. Neonatal seizures in the United States: results of the National Hospital Discharge Survey, 1980–1991. *Neuroepidemiology* 1996; 3: 117–25.
226. Larmay HJ, Strasburger JF. Differential diagnosis and management of the fetus and newborn with an irregular or abnormal heart rate. *Pediatr Clin North Am* 2004; 4: 1033–50.
227. Lawrence MJ, Ford WD, Furness ME et al. Congenital duodenal obstruction: early antenatal ultrasound diagnosis. *Pediatr Surg Int* 2000; 16: 342–5.
228. Le Bihannic A, Beauvais K, Busnel A et al. Prognostic value of EEG in very premature newborns. *Arch Dis Child Fetal Neonatal Ed* 2012; 2: F106–9.
229. Lee EY, Dorkin H, Vargas SO. Congenital pulmonary malformations in pediatric patients: review and update on etiology, classification, and imaging findings. *Radiol Clin North Am* 2011; 5: 921–48.
230. Lee SL, Beyer TD, Kim SS et al. Initial nonoperative management and delayed closure for treatment of giant omphaloceles. *J Pediatr Surg* 2006; 11: 1846–9.
231. Lee TC, Lim FY, Keswani SG et al. Late gestation fetal magnetic resonance imaging-derived total lung volume predicts postnatal survival and need for extracorporeal membrane oxygenation support in isolated congenital diaphragmatic hernia. *J Pediatr Surg* 2011; 6: 1165–71.
232. Lee TW, Skelton RE, Skene C. Routine neonatal examination: effectiveness of trainee paediatrician compared with advanced neonatal nurse practitioner. *Arch Dis Child Fetal Neonatal Ed* 2001; 2: F100–4.
233. Leonard E, Weisman MD, Mohan Pammi MD. Treatment and prevention of bacterial sepsis in the preterm infant. www.uptodate.com/contents/treatment-and-prevention-of-bacterial-sepsis-in-the-preterm-infant.
234. Levinson-Castiel R, Merlob P, Linder N et al. Neonatal abstinence syndrome after in utero exposure to selective serotonin reuptake inhibitors in term infants. *Arch Pediatr Adolesc Med* 2006; 2: 173–6.
235. Levitt MA, Pena A. Outcomes from the correction of anorectal malformations. *Curr Opin Pediatr* 2005; 3: 394–401.
236. Levitt MA, Pena A. Anorectal malformations. *Orphanet J Rare Dis* 2007; 2: 33. Review. Erratum in: *Orphanet J Rare Dis* 2012; 7: 98.
237. Lin PW, Stoll BJ. Necrotising enterocolitis. *Lancet* 2006; 368: 1271–83.
238. Lin ZL, Yu HM, Lin J et al. Mild hypothermia via selective head cooling as neuroprotective therapy in term neonates with perinatal asphyxia: an experience from a single neonatal intensive care unit. *J Perinatol* 2006; 3: 180–4.
239. Lindqvist PG, Erichs K, Molnar C et al. Characteristics and outcome of brachial plexus birth palsy in neonates. *Acta Paediatr* 2012; 6: 579–82.
240. Lippi G, Salvagno GL, Rugolotto S et al. Routine coagulation tests in newborn and young infants. *J Thromb Thrombolysis* 2007; 2: 153–5.
241. Lista G, Fontana P, Castoldi F et al. Does sustained lung inflation at birth improve outcome of preterm infants at risk for respiratory distress syndrome? *Neonatology* 2011; 1: 45–50.
242. Litmanovitz I, Carlo WA. Expectant management of pneumothorax in ventilated neonates. *Pediatrics* 2008; 5: e975–9.
243. Logan JW, Rice HE, Goldberg RN, Cotten CM. Congenital diaphragmatic hernia: a systematic review and summary of best-evidence practice strategies. *J Perinatol* 2007; 9: 535–49.
244. Logan JW, O’Shea TM, Allred EN et al. Early postnatal hypotension and developmental delay at 24 months of age among extremely low gestational age newborns. *Arch Dis Child Fetal Neonatal Ed* 2011; 5: F321–8.
245. Logghe HL, Mason GC, Thornton JG, Stringer MD. A randomized controlled trial of elective preterm delivery of fetuses with gastroschisis. *J Pediatr Surg* 2005; 11: 1726–31.
246. Loh AH, Ong CL, Lam SL et al. Pediatric risk of malignancy index for preoperative evaluation of childhood ovarian tumors. *Pediatr Surg Int* 2012; 3: 259–66.
247. Lokesh L, Kumar P, Murki S, Narang A. A randomized controlled trial of sodium bicarbonate in neonatal resuscitation-effect on immediate outcome. *Resuscitation* 2004; 2: 219–23.
248. Looney CB, Smith JK, Merck LH et al. Intracranial hemorrhage in asymptomatic neonates: prevalence on MR images and relationship to obstetric and neonatal risk factors. *Radiology* 2007; 2: 535–41.
249. Lopez-Gil M, Mantilla I, Blanco T et al. The Size 1 ProSeal laryngeal mask airway in infants: a randomized, noncrossover study with the

- Classic laryngeal mask airway. *Paediatr Anaesth* 2012; 4: 365–70.
250. Lopriore E, Oepkes D, Walther FJ. Neonatal morbidity in twin-twin transfusion syndrome. *Early Hum Dev* 2011; 9: 595–9.
251. Loytved C. Qualitätsbericht 2009. Außerklinische Geburtshilfe in Deutschland. Gesellschaft für Qualität in der außerklinischen Geburtshilfe e. V. (QUAG e. V.) 2010.
252. Loytved C (Hrsg). Außerklinische Geburtshilfe in Deutschland, Qualitätsbericht 2009.
253. Luthy DA, Wardinsky T, Shurtleff DB et al. Cesarean section before the onset of labor and subsequent motor function in infants with meningomyelocele diagnosed antenatally. *N Engl J Med* 1991; 10: 662–6.
254. Maartens IA, Wassenberg T, Buijs J et al. Neurodevelopmental outcome in full-term newborns with refractory neonatal seizures. *Acta Paediatr Scand* 2012; 4: e173–8.
255. Malviya M, Ohlsson A, Shah S. Surgical versus medical treatment with cyclooxygenase inhibitors for symptomatic patent ductus arteriosus in preterm infants. *Cochrane Database Syst Rev* 2008; 1: CD003951.
256. Manganaro R, Mami C, Palmara A et al. Incidence of meconium aspiration syndrome in term meconium-stained babies managed at birth with selective tracheal intubation. *J Perinat Med* 2001; 6: 465–8.
257. Manthey D, Asimiadou S, Stefovskaja V et al. Sulthiame but not levetiracetam exerts neurotoxic effect in the developing rat brain. *Exp Neurol* 2005; 2: 497–503.
258. Martinez-Biarge M, Diez-Sebastian J, Kapellou O et al. Predicting motor outcome and death in term hypoxic-ischemic encephalopathy. *Neurology* 2011; 24: 2055–61.
259. Martins Y, Silva S, Matias A, Blickstein I. Cardiac morbidity in twin-twin transfusion syndrome? *J Perinat Med* 2012; 2: 107–14.
260. Masumoto K, Nagata K, Uesugi T et al. Risk factors for sensorineural hearing loss in survivors with severe congenital diaphragmatic hernia. *Eur J Pediatr* 2007; 6: 607–12.
261. Mayer S, Klaritsch P, Petersen S et al. The correlation between lung volume and liver herniation measurements by fetal MRI in isolated congenital diaphragmatic hernia: a systematic review and meta-analysis of observational studies. *Prenat Diagn* 2011; 11: 1086–96.
262. Mazurier E, Cambonie G, Barbotte E et al. Comparison of chlorpromazine versus morphine hydrochloride for treatment of neonatal abstinence syndrome. *Acta Paediatr Scand* 2008; 10: 1358–61.
263. McIntosh N, Becher JC, Cunningham S et al. Clinical diagnosis of pneumothorax is late: use of trend data and decision support might allow preclinical detection. *Pediatr Res* 2000; 3: 408–15.
264. McNamara PJ, Laique F, Muang-In S, Whyte HE. Milrinone improves oxygenation in neonates with severe persistent pulmonary hypertension of the newborn. *J Crit Care* 2006; 2: 217–22.
265. Mehler K, Wendrich D, Kissgen R et al. Mothers seeing their VLBW infants within 3 h after birth are more likely to establish a secure attachment behavior: evidence of a sensitive period with preterm infants? *J Perinatol* 2011; 6: 404–10.
266. Ment LR, Oh W, Ehrenkranz RA et al. Low-dose indomethacin therapy and extension of intraventricular hemorrhage: a multicenter randomized trial. *J Pediatr* 1994; 6: 951–5.
267. Ment LR, Vohr BR, Makuch RW et al. Prevention of intraventricular hemorrhage by indomethacin in male preterm infants. *J Pediatr* 2004; 6: 832–4.
268. Merei JM, Hutson JM. Embryogenesis of tracheo esophageal anomalies: a review. *Pediatr Surg Int* 2002; 18: 319–26.
269. Metzger R, Wachowiak R, Kluth D. Embryology of the early foregut. *Semin Pediatr Surg* 2011; 3: 136–44.
270. Michaelsson M, Engle MA. Congenital complete heart block: an international study of the natural history. *Cardiovasc Clin* 1972; 3: 85–101.
271. Miller PD, Pollack IF, Pang D, Albright AL. Comparison of simultaneous versus delayed ventriculoperitoneal shunt insertion in children undergoing myelomeningocele repair. *J Child Neurol* 1996; 5: 370–2.
272. Miller SP, Newton N, Ferriero DM et al. Predictors of 30-month outcome after perinatal depression: role of proton MRS and socioeconomic factors. *Pediatr Res* 2002; 1: 71–7.
273. Miller SP, Weiss J, Barnwell A et al. Seizure-associated brain injury in term newborns with perinatal asphyxia. *Neurology* 2002; 4: 542–8.
274. Mizrahi EaK, P. Characterization and classification of neonatal seizures. *Neurology* 1987; 12: 1837–44.

275. Moghal NE, Embleton ND. Management of acute renal failure in the newborn. *Semin Fetal Neonatal Med* 2006; 3: 207–13.
276. Mohamed WA, Ismail M. A randomized, double-blind, placebo-controlled, prospective study of bosentan for the treatment of persistent pulmonary hypertension of the newborn. *J Perinatol* 2012; 8: 608–13.
277. Moise KJ, Jr. Management of rhesus alloimmunization in pregnancy. *Obstet Gynecol* 2002; 3: 600–11.
278. Montassir H, Maegaki Y, Ogura K et al. Associated factors in neonatal hypoglycemic brain injury. *Brain Dev* 2009; 9: 649–56.
279. Moreira-Pinto J, Rocha P, Osorio A et al. Octreotide in the treatment of neonatal postoperative chylothorax: report of three cases and literature review. *Pediatr Surg Int* 2011; 8: 805–9.
280. Morley CJ, Davis PG, Doyle LW et al. Nasal CPAP or intubation at birth for very preterm infants. *N Engl J Med* 2008; 7: 700–8.
281. Mortellaro VE, St Peter SD, Fike FB, Islam S. Review of the evidence on the closure of abdominal wall defects. *Pediatr Surg Int* 2011; 4: 391–7.
282. Moses-Kolko EL, Bogen D, Perel J et al. Neonatal signs after late in utero exposure to serotonin reuptake inhibitors: literature review and implications for clinical applications. *JAMA* 2005; 19: 2372–83.
283. Mosquera C, Miller RS, Simpson LL. Twin-twin transfusion syndrome. *Semin Perinatol* 2012; 3: 182–9.
284. Moya MP, Gow AJ, Califf RM et al. Inhaled ethyl nitrite gas for persistent pulmonary hypertension of the newborn. *Lancet* 2002; 360: 141–3.
285. Mugford M, Elbourne D, Field D. Extracorporeal membrane oxygenation for severe respiratory failure in newborn infants. *Cochrane Database Syst Rev* 2008; 3: CD001340.
286. Müller MJ, Lange M, Paul T, Seeliger S. Stillen unter Methadon- bzw. Buprenorphin-Substitutionstherapie. *Klin Padiatr* 2011; 7: 408–13.
287. Munck A, Gerardin M, Alberti C et al. Clinical outcome of cystic fibrosis presenting with or without meconium ileus: a matched cohort study. *J Pediatr Surg* 2006; 9: 1556–60.
288. Munoz JJ, Mansul AJ, Malpas TJ, Steinbrecher HA. Intrathoracic gastric volvulus mimicking pyloric stenosis. *J Paediatr Child Health* 2003; 2: 149–51.
289. Murray F, Patel HH, Suda RY et al. Expression and activity of cAMP phosphodiesterase isoforms in pulmonary artery smooth muscle cells from patients with pulmonary hypertension: role for PDE1. *Am J Physiol Lung Cell Mol Physiol* 2007; 1: L294–303.
290. Naheed ZJ, Strasburger JF, Deal BJ et al. Fetal tachycardia: mechanisms and predictors of hydrops fetalis. *J Am Coll Cardiol* 1996; 7: 1736–40.
291. Nankervis CA, Giannone PJ, Reber KM. The neonatal intestinal vasculature: contributing factors to necrotizing enterocolitis. *Semin Perinatol* 2008; 2: 83–91.
292. National Collaborating Centre for Women's and Children's Health. *Intrapartum Care: Care of healthy women and their babies during childbirth*. London: RCOG Press 2007; 48–66.
293. Nelson KB, Emery ES. Birth asphyxia and the neonatal brain: what do we know and when do we know it? *Clin Perinatol* 1993; 2: 327–44.
294. Neumann R, Schulzke SM, Bühner C. Oral ibuprofen versus intravenous ibuprofen or intravenous indomethacin for the treatment of patent ductus arteriosus in preterm infants: a systematic review and meta-analysis. *Neonatology* 2012; 1: 9–15.
295. Newburger JW, Keane JF. Intrauterine supraventricular tachycardia. *J Pediatr* 1979; 95: 780–6.
296. Ng E, Taddio A, Ohlsson A. Intravenous midazolam infusion for sedation of infants in the neonatal intensive care unit. *Cochrane Database Syst Rev* 2003; 1: CD002052.
297. Nigrovic LE, Kuppermann N, Macias CG et al. Clinical prediction rule for identifying children with cerebrospinal fluid pleocytosis at very low risk of bacterial meningitis. *JAMA* 2007; 1: 52–60.
298. Nolan JP, Soar J, Zideman DA et al. European Resuscitation Council Guidelines for Resuscitation 2010 Section 1. Executive summary. *Resuscitation* 2010; 10: 1219–76.
299. O'Donnell CP, Kamlin CO, Davis PG et al. Clinical assessment of infant colour at delivery. *Arch Dis Child Fetal Neonatal Ed* 2007; 6: F465–7.
300. Oddie S, Embleton ND. Risk factors for early onset neonatal group B streptococcal sepsis: case-control study. *BMJ* 2002; 7359: 308.
301. Ohlsson A, Walia R, Shah S. Ibuprofen for the treatment of patent ductus arteriosus in preterm

- and/or low birth weight infants. *Cochrane Database Syst Rev* 2005; 4: CD003481.
302. Ohlsson A, Roberts RS, Schmidt B et al. Male/female differences in indomethacin effects in preterm infants. *J Pediatr* 2005; 6: 860–2.
 303. Ohlsson A, Walia R, Shah S. Ibuprofen for the treatment of patent ductus arteriosus in preterm and/or low birth weight infants. *Cochrane Database Syst Rev* 2008; 1: CD003481.
 304. Ohlsson A, Walia R, Shah SS. Ibuprofen for the treatment of patent ductus arteriosus in preterm and/or low birth weight infants. *Cochrane Database Syst Rev* 2010; 4: CD003481.
 305. Ohlsson A, Shah SS. Ibuprofen for the prevention of patent ductus arteriosus in preterm and/or low birth weight infants. *Cochrane Database Syst Rev* 2011; 7: CD004213.
 306. Okamoto T, Takamizawa S, Arai H et al. Esophageal atresia: prognostic classification revisited. *Surgery* 2009; 6: 675–81.
 307. Olesevich M, Alexander F, Khan M, Cotman K. Gastroschisis revisited: role of intraoperative measurement of abdominal pressure. *J Pediatr Surg* 2005; 5: 789–92.
 308. Olsen O, Jewell MD. Home versus hospital birth. *Cochrane Database Syst Rev* 2000; 2: CD000352.
 309. Opdal SH, Rognum TO. Gene variants predisposing to SIDS: current knowledge. *Forensic Sci Med Pathol* 2011; 1: 26–36.
 310. Oppenheimer C. Organising midwifery led care in The Netherlands. *BMJ* 1993; 307: 1400–2.
 311. Osborn DA, Jeffery HE, Cole MJ. Opiate treatment for opiate withdrawal in newborn infants. *Cochrane Database Syst Rev* 2010; 10: CD002059.
 312. Ostrea EM, Villanueva-Uy ET, Natarajan G, Uy HG. Persistent pulmonary hypertension of the newborn: pathogenesis, etiology, and management. *Paediatr Drugs* 2006; 3: 179–88.
 313. Owen A, Marven S, Johnson P et al. Gastroschisis: a national cohort study to describe contemporary surgical strategies and outcomes. *J Pediatr Surg* 2010; 9: 1808–16.
 314. Ozek E, Soll R, Schimmel MS. Partial exchange transfusion to prevent neurodevelopmental disability in infants with polycythemia. *Cochrane Database Syst Rev* 2010; 1: CD005089.
 315. Painter MJ, Scher MS, Stein AD et al. Phenobarbital compared with phenytoin for the treatment of neonatal seizures. *N Engl J Med* 1999; 7: 485–9.
 316. Pandey R, Jackson M, Ajab S et al. Subclavian flap repair: review of 399 patients at median follow-up of fourteen years. *Ann Thorac Surg* 2006; 4: 1420–8.
 317. Pang JW, Heffelfinger JD, Huang GJ et al. Outcomes of planned home births in Washington State: 1989–1996. *Obstet Gynecol* 2002; 2: 253–9.
 318. Park MD (ed). *Park's Pediatric Cardiology for Practitioners*. 6th ed. St. Louis: Mosby 2014.
 319. Park MK. *Pediatric cardiology for practitioners*. St. Louis: Mosby 2008.
 320. Pastor AC, Phillips JD, Fenton SJ et al. Routine use of a SILASTIC spring-loaded silo for infants with gastroschisis: a multicenter randomized controlled trial. *J Pediatr Surg* 2008; 10: 1807–12.
 321. Pearson DL, Dawling S, Walsh WF et al. Neonatal pulmonary hypertension – urea-cycle intermediates, nitric oxide production, and carbamoyl-phosphate synthetase function. *N Engl J Med* 2001; 24: 1832–8.
 322. *Pediatrics AAO. Red Book: Amer Assn of Pediatrics* 2009.
 323. Pena A, Hong A. Advances in the management of anorectal malformations. *Am J Surg* 2000; 5: 370–6.
 324. Perlman JM, Risser R. Can asphyxiated infants at risk for neonatal seizures be rapidly identified by current high-risk markers? *Pediatrics* 1996; 4: 456–62.
 325. Perlman JM, Wyllie J, Kattwinkel J et al. Neonatal Resuscitation Chapter C. Part 11: Neonatal resuscitation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation* 2010; 16 Suppl 2: S516–38. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
 326. Phibbs RH. Hemolytic Disease of the Newborn. In: Rudolph AM (Hrsg). *Rudolph's Pediatrics*. Stamford: Appleton & Lange 1996; 1193–200.
 327. Pickering LK. *Red Book: 2006 Report of the committee on infectious diseases: American Academy of Pediatrics* 2006. Siehe Redbook Updates.
 328. Pini-Prato A, Martucciello G, Jasonni V. Rectal suction biopsy in the diagnosis of intestinal dysganglionoses: 5-year experience with Solo-RBT in 389 patients. *J Pediatr Surg* 2006; 6: 1043–8.
 329. Poets A, Steinfeldt R, Poets CF. Sudden deaths and severe apparent life-threatening events in

- term infants within 24 hours of birth. *Pediatrics* 2011; 4: e869–73.
330. Kurz R, Kenner T, Poets C et al. (Hrsg). *Der Plötzliche Säuglingstod. Grundlagen, Risikofaktoren, Prävention, Elternberatung*. 2. Aufl. Wien: Springer 2014.
331. Poets CF, Bacher M. Treatment of upper airway obstruction and feeding problems in Robin-like phenotype. *J Pediatr* 2011; 6: 887–92.
332. Poets CF. Der plötzliche Kindstod. In: Lentze MJ, Schaub J, Schulte FJ, Spranger J (Hrsg). *Pädiatrie – Grundlagen und Praxis*. Heidelberg: Springer 2012.
333. Poki HO, Holland AJ, Pitkin J. Double bubble, double trouble. *Pediatr Surg Int* 2005; 6: 428–31.
334. Printzlau A, Andersen M. Pierre Robin sequence in Denmark: a retrospective population-based epidemiological study. *Cleft Palate Craniofac J* 2004; 1: 47–52.
335. Rabe H, Reynolds G, Diaz-Rossello J. A systematic review and meta-analysis of a brief delay in clamping the umbilical cord of preterm infants. *Neonatology* 2008; 2: 138–44.
336. Ralston M, Hazinski MF, Zaritsky AL et al. *PALS Provider Manual*. Dallas: American Heart Association and American Academy of Pediatrics 2006.
337. Rao P. Neonatal gastrointestinal imaging. *Eur J Radiol* 2006; 2: 171–86.
338. Rao SC, Basani L, Simmer K et al. Peritoneal drainage versus laparotomy as initial surgical treatment for perforated necrotizing enterocolitis or spontaneous intestinal perforation in preterm low birth weight infants. *Cochrane Database Syst Rev* 2011; 6: CD006182.
339. Rees CM, Eaton S, Kiely EM et al. Peritoneal drainage or laparotomy for neonatal bowel perforation? A randomized controlled trial. *Ann Surg* 2008; 1: 44–51.
340. Rennie JM, Cooke RW. Prolonged low dose indomethacin for persistent ductus arteriosus of prematurity. *Arch Dis Child* 1991; 1: 55–8.
341. Rennie JM. Neonatal seizures. *Eur J Pediatr* 1997; 2: 83–7.
342. Rennie JM, Chorley G, Boylan GB et al. Non-expert use of the cerebral function monitor for neonatal seizure detection. *Arch Dis Child Fetal Neonatal Ed* 2004; 1: F37–40.
343. Rennie JM. *Robertson's Textbook of Neonatology*. Edinburgh: Churchill Livingstone 2005.
344. Rennie JM. *Rennie & Robertson's Textbook of Neonatology*. Edinburgh: Churchill Livingstone 2012.
345. Richmond S, Wyllie J. *European Resuscitation Council Guidelines for Resuscitation 2010 Section 7. Resuscitation of babies at birth*. *Resuscitation* 2010; 10: 1389–99. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
346. Roberts D, Neilson JP, Kilby M, Gates S. Interventions for the treatment of twin-twin transfusion syndrome. *Cochrane Database Syst Rev* 2008; 1: CD002073.
347. Robertson C, Finer N. Term infants with hypoxic-ischemic encephalopathy: outcome at 3.5 years. *Dev Med Child Neurol* 1985; 4: 473–84.
348. Roehr CC, Hansmann G, Hoehn T, Bühner C. The 2010 Guidelines on Neonatal Resuscitation (AHA, ERC, ILCOR): similarities and differences – what progress has been made since 2005? *Klin Padiatr* 2011; 5: 299–307. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
349. Rooks VJ, Eaton JP, Ruess L et al. Prevalence and evolution of intracranial hemorrhage in asymptomatic term infants. *AJNR Am J Neuroradiol* 2008; 6: 1082–9.
350. Rosenberg AA. Outcome in term infants treated with inhaled nitric oxide. *J Pediatr* 2002; 3: 284–7.
351. Rowland CA, Correa A, Cragan JD, Alverson CJ. Are encephaloceles neural tube defects? *Pediatrics* 2006; 3: 916–23.
352. Ruano R, Yoshisaki CT, da Silva MM et al. A randomized controlled trial of fetal endoscopic tracheal occlusion versus postnatal management of severe isolated congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2012; 1: 20–7.
353. Rutherford M, Ramenghi LA, Edwards AD et al. Assessment of brain tissue injury after moderate hypothermia in neonates with hypoxic-ischaemic encephalopathy: a nested substudy of a randomised controlled trial. *Lancet Neurol* 2010; 1: 39–45.
354. Rutherford MA, Pennock JM, Counsell SJ et al. Abnormal magnetic resonance signal in the internal capsule predicts poor neurodevelopmental outcome in infants with hypoxic-ischemic encephalopathy. *Pediatrics* 1998; 102: 323–8.

355. Ruttenstock E, Puri P. Systematic review and meta-analysis of enterocolitis after one-stage transanal pull-through procedure for Hirschsprung's disease. *Pediatr Surg Int* 2010; 11: 1101–5.
356. Sallmon H, Weber SC, von Gise A et al. Ductal closure in neonates: a developmental perspective on platelet-endothelial interactions. *Blood Coagul Fibrinolysis* 2011; 3: 242–4.
357. Sallmon H, Sola-Visner M. Clinical and research issues in neonatal anemia and thrombocytopenia. *Curr Opin Pediatr* 2012; 1: 16–22.
358. Sankar R, Painter MJ. Neonatal seizures: after all these years we still love what doesn't work. *Neurology* 2005; 5: 776–7.
359. Schermuly RT, Pullamsetti SS, Kwapiszewska G et al. Phosphodiesterase 1 upregulation in pulmonary arterial hypertension: target for reverse-remodeling therapy. *Circulation* 2007; 17: 2331–9.
360. Schmidt B, Davis P, Moddemann D et al. Long-term effects of indomethacin prophylaxis in extremely-low-birth-weight infants. *N Engl J Med* 2001; 26: 1966–72.
361. Schmidt B, Roberts RS, Fanaroff A et al. Indomethacin prophylaxis, patent ductus arteriosus, and the risk of bronchopulmonary dysplasia: further analyses from the Trial of Indomethacin Prophylaxis in Preterms (TIPP). *J Pediatr* 2006; 6: 730–4.
362. Schnabl KL, Van Aerde JE, Thomson AB, Clandinin MT. Necrotizing enterocolitis: a multifactorial disease with no cure. *World J Gastroenterol* 2008; 14: 2142–61.
363. Schorscher-Petcu A, Sotocinal S, Ciura S et al. Oxytocin-induced analgesia and scratching are mediated by the vasopressin-1A receptor in the mouse. *J Neurosci* 2010; 24: 8274–84.
364. Schrag SJ, Zywicki S, Farley MM et al. Group B streptococcal disease in the era of intrapartum antibiotic prophylaxis. *N Engl J Med* 2000; 1: 15–20.
365. Schrag SJ, Zell ER, Lynfield R et al. A population-based comparison of strategies to prevent early-onset group B streptococcal disease in neonates. *N Engl J Med* 2002; 4: 233–9.
366. Schrag SJ, Stoll BJ. Early-onset neonatal sepsis in the era of widespread intrapartum chemoprophylaxis. *Pediatr Infect Dis J* 2006; 10: 939–40.
367. Shah DK, Boylan GB, Rennie JM. Monitoring of seizures in the newborn. *Arch Dis Child Fetal Neonatal Ed* 2012; 1: F65–9.
368. Shah P, Ohlsson A. Sildenafil for pulmonary hypertension in neonates. *Cochrane Database Syst Rev* 2011; 8: CD005494.
369. Shah PS, Hellmann J, Adatia I. Clinical characteristics and follow up of Down syndrome infants without congenital heart disease who presented with persistent pulmonary hypertension of newborn. *J Perinat Med* 2004; 2: 168–70.
370. Shah PS, Hellmann J, Adatia I. Clinical characteristics and follow up of Down syndrome infants without congenital heart disease who presented with persistent pulmonary hypertension of newborn. *J Perinat Med* 2004; 2: 168–70.
371. Shankaran S, Laptook AR, Ehrenkranz RA et al. Whole-body hypothermia for neonates with hypoxic-ischemic encephalopathy. *N Engl J Med* 2005; 15: 1574–84.
372. Shankaran S, Langer JC, Kazzi SN et al. Cumulative index of exposure to hypocarbia and hyperoxia as risk factors for periventricular leukomalacia in low birth weight infants. *Pediatrics* 2006; 4: 1654–9.
373. Sharma R, Tepas JJ, Hudak ML et al. Neonatal gut injury and infection rate: impact of surgical debridement on outcome. *Pediatr Surg Int* 2005; 12: 977–82.
374. Shaw-Smith C. Oesophageal atresia, tracheo-oesophageal fistula, and the VACTERL association: review of genetics and epidemiology. *J Med Genet* 2006; 7: 545–54.
375. Shawis R, Antao B. Prenatal bowel dilatation and the subsequent postnatal management. *Early Hum Dev* 2006; 5: 297–303.
376. Siegfried N, van der Merwe L, Brocklehurst P, Sint TT. Antiretrovirals for reducing the risk of mother-to-child transmission of HIV infection. *Cochrane Database Syst Rev* 2011; 7: CD003510.
377. Silberbach M, Hannon D. Presentation of congenital heart disease in the neonate and young infant. *Pediatr Rev* 2007; 4: 123–31.
378. Simonneau G, Gatzoulis MA, Adatia I et al. Updated clinical classification of pulmonary hypertension. *J Am Coll Cardiol* 2011; 62: D34–41.
379. Simpson J, Yates RW, Sharland GK. Irregular heart rate in the fetus – not always benign. *Cardiol Young* 1996; 6: 28–31.
380. Simpson JM. Fetal arrhythmias. *Ultrasound Obstet Gynecol* 2006; 6: 599–606.
381. Singh M, Owen A, Gull S et al. Surgery for intestinal perforation in preterm neonates: anastomosis vs stoma. *J Pediatr Surg* 2006; 4: 725–9.

382. Singh RR, Warren PS, Reece TB et al. Early repair of complete atrioventricular septal defect is safe and effective. *Ann Thorac Surg* 2006; 5: 1598–602.
383. Skinner J. Diagnosis of patent ductus arteriosus. *Semin Neonatol* 2001; 1: 49–61.
384. Slaghekke F, Kist WJ, Oepkes D et al. Twin anemia-polycythemia sequence: diagnostic criteria, classification, perinatal management and outcome. *Fetal Diagn Ther* 2010; 4: 181–90.
385. El Shahed Al, Dargaville PA, Ohlsson A, Sool R. Surfactant for meconium aspiration syndrome in full term infants. *Cochrane Database Syst Rev* 2014; 12: CD002054.
386. Southall DP, Richards J, Hardwick RA et al. Prospective study of fetal heart rate and rhythm patterns. *Arch Dis Child* 1980; 7: 506–11.
387. Spitz L. Esophageal atresia. Lessons I have learned in a 40-year experience. *J Pediatr Surg* 2006; 10: 1635–40.
388. Spitzmiller RE, Phillips T, Meinzen-Derr J, Hoath SB. Amplitude-integrated EEG is useful in predicting neurodevelopmental outcome in full-term infants with hypoxic-ischemic encephalopathy: a meta-analysis. *J Child Neurol* 2007; 9: 1069–78.
389. Statistiek Centraal Bureau. Length and Weight at Birth and Breastfeeding (Bevalling, lengte en gewicht bij geboorte en borstvoeding). *Delivery* 2008.
390. Stephenson JM, Du JN, Oliver TK Jr. The effect of cooling on blood gas tensions in newborn infants. *J Pediatr* 1970; 6: 848–52.
391. Stockman JA. Overview of the state of the art of Rh disease: history, current clinical management, and recent progress. *J Pediatr Hematol Oncol* 2001; 6: 385–93.
392. Stoll BJ, Hansen N, Fanaroff AA et al. Changes in pathogens causing early-onset sepsis in very-low-birth-weight infants. *N Engl J Med* 2002; 4: 240–7.
393. Stoll C, Alembik Y, Dott B, Roth MP. Associated malformations in patients with anorectal anomalies. *Eur J Med Genet* 2007; 4: 281–90.
394. Strasburger JF. Fetal arrhythmias. *Prog Pediatr Cardiol* 2000; 1: 1–17.
395. Strasburger JF, Cuneo BF, Michon MM et al. Amiodarone therapy for drug-refractory fetal tachycardia. *Circulation* 2004; 3: 375–9.
396. Strasburger JF. Prenatal diagnosis of fetal arrhythmias. *Clin Perinatol* 2005; 4: 891–912.
397. Strasburger JF, Wakai RT. Fetal cardiac arrhythmia detection and in utero therapy. *Nat Rev Cardiol* 2010; 5: 277–90.
398. Sudarshan CD, Cochrane AD, Jun ZH et al. Repair of coarctation of the aorta in infants weighing less than 2 kilograms. *Ann Thorac Surg* 2006; 1: 158–63.
399. Sulzbacher S, Farwell JR, Temkin N et al. Late cognitive effects of early treatment with phenobarbital. *Clin Pediatr (Phila)* 1999; 7: 387–94.
400. Synnes AR, Chien LY, Peliowski A et al. Variations in intraventricular hemorrhage incidence rates among Canadian neonatal intensive care units. *J Pediatr* 2001; 4: 525–31.
401. Tan BH, Pundi KN, Van Norstrand DW et al. Sudden infant death syndrome-associated mutations in the sodium channel beta subunits. *Heart Rhythm* 2010; 6: 771–8.
402. Te Pas AB, Walther FJ. A randomized, controlled trial of delivery-room respiratory management in very preterm infants. *Pediatrics* 2007; 2: 322–9.
403. Teng RJ, Wu TJ. Persistent pulmonary hypertension of the newborn. *J Formos Med Assoc* 2013; 4: 177–84.
404. Tharp BR. Neonatal seizures and syndromes. *Epilepsia* 2002; 43 Suppl 3: 2–10.
405. Thebaud B, Michelakis ED, Wu XC et al. Oxygen-sensitive Kv channel gene transfer confers oxygen responsiveness to preterm rabbit and remodeled human ductus arteriosus: implications for infants with patent ductus arteriosus. *Circulation* 2004; 11: 1372–9.
406. Toet MC, van der Meij W, de Vries LS et al. Comparison between simultaneously recorded amplitude integrated electroencephalogram (cerebral function monitor) and standard electroencephalogram in neonates. *Pediatrics* 2002; 5: 772–9.
407. Townsend J, Wolke D, Hayes J et al. Routine examination of the newborn: the EMREN study. Evaluation of an extension of the midwife role including a randomised controlled trial of appropriately trained midwives and paediatric senior house officers. *Health Technol Assess* 2004; 14: iii-iv, ix-xi, 1–100.
408. Trappe HJ. Acute therapy of maternal and fetal arrhythmias during pregnancy. *J Intensive Care Med* 2006; 5: 305–15.
409. Turnpenny PE, S. Emery's Elements of Medical Genetics. Amsterdam: Elsevier 2012.

410. Tworetzky W, Bristow J, Moore P et al. Inhaled nitric oxide in neonates with persistent pulmonary hypertension. *Lancet* 2001; 357: 118–20.
411. Tyson JE, Parikh NA, Langer J et al. National Institute of Child Health, Human Development Neonatal Research N. Intensive care for extreme prematurity – moving beyond gestational age. *N Engl J Med* 2008; 16: 1672–81.
412. Vain NE, Szlyd EG, Prudent LM et al. Oropharyngeal and nasopharyngeal suctioning of meconium-stained neonates before delivery of their shoulders: multicentre, randomised controlled trial. *Lancet* 2004; 364: 597–602.
413. Valsky DV, Eixarch E, Martinez JM, Gratacos E. Selective intrauterine growth restriction in monochorionic diamniotic twin pregnancies. *Prenat Diagn* 2010; 8: 719–26.
414. Van den Hout L, Tibboel D, Vijfhuizen S et al. The VICI-trial: high frequency oscillation versus conventional mechanical ventilation in newborns with congenital diaphragmatic hernia: an international multicentre randomized controlled trial. *BMC Pediatr* 2011; 11: 98.
415. Van Eijck FC, Hoogeveen YL, van Weel C et al. Minor and giant omphalocele: long-term outcomes and quality of life. *J Pediatr Surg* 2009; 7: 1355–9.
416. Van Overmeire B, Smets K, Lecoutere D et al. A comparison of ibuprofen and indomethacin for closure of patent ductus arteriosus. *N Engl J Med* 2000; 10: 674–81.
417. Van Praagh R, Perez-Trevino C, Lopez-Cuellar M et al. Transposition of the great arteries with posterior aorta, anterior pulmonary artery, subpulmonary conus and fibrous continuity between aortic and atrioventricular valves. *Am J Cardiol* 1971; 6: 621–31.
418. Vanhaesebrouck S, Zonnenberg I, Vandervoort P et al. Conservative treatment for patent ductus arteriosus in the preterm. *Arch Dis Child Fetal Neonatal Ed* 2007; 4: F244–7.
419. Vannucci RC, Perlman JM. Interventions for perinatal hypoxic-ischemic encephalopathy. *Pediatrics* 1997; 6: 1004–14.
420. Vannucci RC, Vannucci SJ. Hypoglycemic brain injury. *Sem Neonatol* 2001; 2: 147–55.
421. Vasta I, Kinali M, Messina S et al. Can clinical signs identify newborns with neuromuscular disorders? *J Pediatr* 2005; 1: 73–9.
422. Vegunta RK, Wallace LJ, Leonardi MR et al. Perinatal management of gastroschisis: analysis of a newly established clinical pathway. *J Pediatr Surg* 2005; 3: 528–34.
423. Velez M, Jansson LM. The Opioid dependent mother and newborn dyad: non-pharmacologic care. *J Addict Med* 2008; 3: 113–20.
424. Vento M, Moro M, Escrig R et al. Preterm resuscitation with low oxygen causes less oxidative stress, inflammation, and chronic lung disease. *Pediatrics* 2009; 3: e439–49.
425. Verbeek RJ, Heep A, Maurits NM et al. Fetal endoscopic myelomeningocele closure preserves segmental neurological function. *Dev Med Child Neurol* 2012; 1: 15–22.
426. Vermold HT, Kitterman JA, Phibbs RH et al. Aortic blood pressure during the first 12 hours of life in infants with birth weight 610 to 4,220 grams. *Pediatrics* 1981; 5: 607–13.
427. Volpe JJ. *Neurology of the newborn*. 5th ed. Philadelphia: W. B. Saunders 2009.
428. Waldman JD, Wernly JA. Cyanotic congenital heart disease with decreased pulmonary blood flow in children. *Pediatr Clin North Am* 1999; 2: 385–404.
429. Walker GM, Raine PA. Biliious vomiting in the newborn: how often is further investigation undertaken? *J Pediatr Surg* 2007; 4: 714–6.
430. Walsh P. *Cardiac Arrhythmias*, in *Pediatric Cardiology*. Amsterdam: Elsevier 2007.
431. Walsh-Sukys MC, Tyson JE, Wright LL et al. Persistent pulmonary hypertension of the newborn in the era before nitric oxide: practice variation and outcomes. *Pediatrics* 2000; 105: 14–20.
432. Watkinson M, Tiron I. Events before the diagnosis of a pneumothorax in ventilated neonates. *Arch Dis Child Fetal Neonatal Ed* 2001; 3: F201–3.
433. Wedgwood S, McMullan DM, Bekker JM et al. Role for endothelin-1-induced superoxide and peroxynitrite production in rebound pulmonary hypertension associated with inhaled nitric oxide therapy. *Circ Res* 2001; 4: 357–64.
434. Weijerman ME, van Furth AM, van der Mooren MD et al. Prevalence of congenital heart defects and persistent pulmonary hypertension of the neonate with Down syndrome. *Eur J Pediatr* 2010; 10: 1195–9.
435. Steinhorn R. Diagnosis and treatment of pulmonary hypertension in infancy. *Early Hum Dev* 2013; 89: 865–74.
436. Wellmann S, Benzing J, Cippa G et al. High co-peptin concentrations in umbilical cord blood

- after vaginal delivery and birth acidosis. *J Clin Endocrinol Metab* 2010; 11: 5091–6.
437. Wilhelm L, Borgers H. The δ -equal sign: a novel marker in the diagnosis of fetal isolated cleft palate. *Ultrasound Obstet Gynecol* 2010; 4: 439–44.
438. Williams H. Green for danger! Intestinal malrotation and volvulus. *Arch Dis Child Educ Pract Ed* 2007; 3: 87–91.
439. Wirrell EC. Neonatal seizures: to treat or not to treat? *Semin Pediatr Neurol* 2005; 2: 97–105.
440. Wiswell TE, Baumgart S, Gannon CM, Spitzer AR. No lumbar puncture in the evaluation for early neonatal sepsis: will meningitis be missed? *Pediatrics* 1995; 6: 803–6.
441. Wiswell TE. Handling the meconium-stained infant. *Semin Neonatol* 2001; 3: 225–31.
442. Wiswell TE, Knight GR, Finer NN et al. A multicenter, randomized, controlled trial comparing Surfactin (Lucinactant) lavage with standard care for treatment of meconium aspiration syndrome. *Pediatrics* 2002; 6: 1081–7.
443. Wiswell TE, Tin W, Ohler K. Evidence-based use of adjunctive therapies to ventilation. *Clin Perinatol* 2007; 1: 191–204.
444. Woodgate PG, Davies MW. Permissive hypercapnia for the prevention of morbidity and mortality in mechanically ventilated newborn infants. *Cochrane Database Syst Rev* 2001; 2: CD002061.
445. World Health Organization. Hypoglycaemia of the newborn. Review of the literature. Geneva 1997.
446. Wyllie J, Perlman JM, Kattwinkel J et al. Part 11: Neonatal resuscitation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2010; 81 Suppl 1: e260–87. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
447. Xie HG, Cao YJ, Gauda EB et al. Clonidine clearance matures rapidly during the early postnatal period: a population pharmacokinetic analysis in newborns with neonatal abstinence syndrome. *J Clin Pharmacol* 2011; 4: 502–11.
448. Yinon Y, Grisaru-Granovsky S, Chaddha V et al. Perinatal outcome following fetal chest shunt insertion for pleural effusion. *Ultrasound Obstet Gynecol* 2010; 1: 58–64.
449. Young TE, Magnum B. Neofax 2009. Montvale, NJ: Thomson Reuter 2009.
450. Zhao L, Mason NA, Morrell NW et al. Sildenafil inhibits hypoxia-induced pulmonary hypertension. *Circulation* 2001; 4: 424–8.
451. Zimmermann-Baer U, Notzli U, Rentsch K, Bucher HU. Finnegan neonatal abstinence scoring system: normal values for first 3 days and weeks 5–6 in non-addicted infants. *Addiction* 2010; 3: 524–8.
452. Zola EM, Gunkel JH, Chan RK et al. Comparison of three dosing procedures for administration of bovine surfactant to neonates with respiratory distress syndrome. *J Pediatr* 1993; 3: 453–9.

Teil IV Spezielle Notfälle und Probleme im Neugeborenen- und jungen Säuglingsalter

1. AWMF-Leitlinie 027/016 (www.awmf.org). Diagnostik und Therapieansätze bei Mitochondriopathien im Kindes- und Jugendalter.
2. Allen D, Verjee S, Rees S et al. Platelet transfusion in neonatal alloimmune thrombocytopenia. *Blood* 2007; 1: 388–9.
3. Altirkawi K, Rozycki HJ. Hypocalcemia is common in the first 48 h of life in ELBW infants. *J Perinat Med* 2008; 4: 348–53.
4. Andrew M, Vegh P, Caco C et al. A randomized, controlled trial of platelet transfusions in thrombocytopenic premature infants. *J Pediatr* 1993; 2: 285–91.
5. Arnold DM, Smith JW, Kelton JG. Diagnosis and management of neonatal alloimmune thrombocytopenia. *Transfus Med Rev* 2008; 4: 255–67.
6. Beardsley DS. Venous thromboembolism in the neonatal period. *Semin Perinatol* 2007; 4: 250–3.
7. Bhat MA, Charoo BA, Bhat JI et al. Magnesium sulfate in severe perinatal asphyxia: a randomized, placebo-controlled trial. *Pediatrics* 2009; 5: e764–9.
8. Bhatia J. Fluid and electrolyte management in the very low birth weight neonate. *J Perinatol* 2006; 26 Suppl 1: S19–21.
9. Bosenberg AT, Brown RA. Management of congenital diaphragmatic hernia. *Curr Opin Anaesthesiol* 2008; 3: 323–31.
10. Broemling N, Campbell F. Anesthetic management of congenital tracheoesophageal fistula. *Paediatr Anaesth* 2011; 11: 1092–9.

11. Bussel JB, Sola-Visner M. Current approaches to the evaluation and management of the fetus and neonate with immune thrombocytopenia. *Semin Perinatol* 2009; 1: 35–42.
12. Bussel JB, Buchanan GR, Nugent DJ et al. A randomized, double-blind study of romiplostim to determine its safety and efficacy in children with immune thrombocytopenia. *Blood* 2011; 1: 28–36.
13. Chalmers EA. Perinatal stroke – risk factors and management. *Br J Haematol* 2005; 3: 333–43.
14. Chowdhury O, Greenough A. Neonatal ventilatory techniques – which are best for infants born at term? *Arch Med Sci* 2011; 3: 381–7.
15. Coombs CJ, Richardson PW, Dowling GJ et al. Brachial artery thrombosis in infants: an algorithm for limb salvage. *Plast Reconstr Surg* 2006; 5: 1481–8.
16. Cotton R, Suarez S, Reese J. Unexpected extrarenal effects of loop diuretics in the preterm neonate. *Acta Paediatr* 2012; 101: 835–45.
17. Gentili A, Giuntoli L, Bacchi Reggiani ML et al. Neonatal congenital diaphragmatic hernia: respiratory and blood-gas derived indices in choosing surgical timing. *Minerva Anestesiol* 2012; 10: 1117–25.
18. Grammatikopoulos T, Greenough A, Pallidis C, Davenport M. Benefits and risks of calcium resonant therapy in hyperkalaemic preterm infants. *Acta Paediatr* 2003; 1: 118–20.
19. Greenway A, Massicotte MP, Monagle P. Neonatal thrombosis and its treatment. *Blood Rev* 2004; 2: 75–84.
20. Gupta RK, Parelkar SV, Oak SN et al. Early experience with thoracoscopic repair of congenital diaphragmatic hernias in pediatric age group: results and lessons learned. *Pediatr Surg Int* 2011; 6: 563–6.
21. Hardman JG, Wills JS. The development of hypoxaemia during apnoea in children: a computational modelling investigation. *Br J Anaesth* 2006; 4: 564–70.
22. Holland AJ, Walker K, Badawi N. Gastroschisis: an update. *Pediatr Surg Int* 2010; 9: 871–8.
23. Jain A, Agarwal R, Sankar MJ et al. Hypocalcemia in the newborn. *Indian J Pediatr* 2010; 10: 1123–8.
24. Johnson LH, Mapp DC, Rouse DJ et al. Association of cord blood magnesium concentration and neonatal resuscitation. *J Pediatr* 2012; 4: 573–7.
25. Johr M, Berger TM. Fiberoptic intubation through the laryngeal mask airway (LMA) as a standardized procedure. *Paediatr Anaesth* 2004; 7: 614.
26. Kiefel V, Bassler D, Kroll H et al. Antigen-positive platelet transfusion in neonatal allo-immune thrombocytopenia (NAIT). *Blood* 2006; 9: 3761–3.
27. Kim JH, Lee YS, Kim SH et al. Does umbilical vein catheterization lead to portal venous thrombosis? Prospective US evaluation in 100 neonates. *Radiology* 2001; 3: 645–50.
28. King RI, Mackay RJ, Florkowski CM, Lynn AM. Electrolytes in sick neonates – which sodium is the right answer? *Arch Dis Child Fetal Neonatal Ed* 2013; 98: F74–6.
29. Laing IA, Wong CM. Hypernatraemia in the first few days: is the incidence rising? *Arch Dis Child Fetal Neonatal Ed* 2002; 3: F158–62.
30. Lau KK, Stoffman JM, Williams S et al. Neonatal renal vein thrombosis: review of the English-language literature between 1992 and 2006. *Pediatrics* 2007; 5: e1278–84.
31. Lietman SA, Germain-Lee EL, Levine MA. Hypercalcemia in children and adolescents. *Curr Opin Pediatr* 2010; 4: 508–15.
32. Lorenz JM, Kleinman LI, Markarian K. Potassium metabolism in extremely low birth weight infants in the first week of life. *J Pediatr* 1997; 131: 81–6.
33. MacDonald MGM, Martha D, Seshia Mary M. *Avery's Neonatology: Pathophysiology and Management of the Newborn*. Lippincott Williams & Wilkins (LWW) 2005.
34. Malowany JI, Monagle P, Knoppert DC et al. Enoxaparin for neonatal thrombosis: a call for a higher dose for neonates. *Thromb Res* 2008; 6: 826–30.
35. Manco-Johnson MJ, Grabowski EF, Hellgreen M et al. Recommendations for tPA thrombolysis in children. On behalf of the Scientific Subcommittee on Perinatal and Pediatric Thrombosis of the Scientific and Standardization Committee of the International Society of Thrombosis and Haemostasis. *Thromb Haemost* 2002; 1: 157–8.
36. Mildenerger E, Versmold HT. Pathogenesis and therapy of non-oliguric hyperkalaemia of the premature infant. *Eur J Pediatr* 2002; 8: 415–22.
37. Mitsiakos G, Papaioannou G, Giougi E et al. Is the use of rFVIIa safe and effective in bleed-

- ing neonates? A retrospective series of 8 cases. *J Pediatr Hematol Oncol* 2007; 3: 145–50.
38. Murray N BS, Casbard A, Murphy M et al. PlaNeT Study Grp. A multicentre prospective observational study of platelet transfusion practice in neonates with severe thrombocytopenia. 48th Annual Meeting of the American-Society-of-Hematology 2006; 287A.
 39. Murray NA, Howarth LJ, McCloy MP et al. Platelet transfusion in the management of severe thrombocytopenia in neonatal intensive care unit patients. *Transfus Med* 2002; 1: 35–41.
 40. Noone D, Kieran E, Molloy EJ. Serum Magnesium in the First Week of Life in Extremely Low Birth Weight Infants. *Neonatology* 2012; 4: 274–7.
 41. Nowak-Gottl U, von Kries R, Gobel U. Neonatal symptomatic thromboembolism in Germany: two year survey. *Arch Dis Child Fetal Neonatal Ed* 1997; 3: F163–7.
 42. O'Grady NP, Alexander M, Dellinger EP et al. Guidelines for the prevention of intravascular catheter-related infections. The Hospital Infection Control Practices Advisory Committee, Center for Disease Control and Prevention, U. S. *Pediatrics* 2002; 5: e51.
 43. Orkin SH, Nathan D, Ginsburg D. Nathan and Oski's Hematology of Infancy and Childhood. Saunders W B Co 2008.
 44. Prietsch V, Lindner M, Zschocke J et al. Emergency management of inherited metabolic diseases. *J Inherit Metab Dis* 2002; 7: 531–46.
 45. Reiss I, Schaible T, van den Hout L et al. Standardized postnatal management of infants with congenital diaphragmatic hernia in Europe: the CDH EURO Consortium consensus. *Neonatology* 2010; 4: 354–64.
 46. Rivers A, Slayton WB. Congenital cytopenias and bone marrow failure syndromes. *Semin Perinatol* 2009; 1: 20–8.
 47. Roberts I, Stanworth S, Murray NA. Thrombocytopenia in the neonate. *Blood Rev* 2008; 4: 173–86.
 48. Rouse DJ, Hirtz DG, Thom E et al. A randomized, controlled trial of magnesium sulfate for the prevention of cerebral palsy. *New Engl J Med* 2008; 9: 895–905.
 49. Sallmon H, Gutti RK, Ferrer-Marin F et al. Increasing platelets without transfusion: is it time to introduce novel thrombopoietic agents in neonatal care? *J Perinatol* 2010; 12: 765–9.
 50. Sallmon H, Sola-Visner M. Clinical and research issues in neonatal anemia and thrombocytopenia. *Curr Opin Pediatr* 2012; 1: 16–22.
 51. Saudubray JM, Sedel F, Walter JH. Clinical approach to treatable inborn metabolic diseases: an introduction. *J Inherit Metab Dis* 2006; 2–3: 261–74.
 52. Saxonhouse MA, Manco-Johnson MJ. The evaluation and management of neonatal coagulation disorders. *Semin Perinatol* 2009; 1: 52–65.
 53. Schmidt B, Andrew M. Neonatal thrombosis: report of a prospective Canadian and international registry. *Pediatrics* 1995; 96: 939–43.
 54. Schmidt J SJ, Becke K, Giest J, Schmitz B. Handlungsempfehlung zur Rapid-Sequence-Induction im Kindesalter. *Anästh Intensivmed* 2007; 48: S88–93.
 55. Scriver CR, Beadet AL, Sly WS, Valle D. Metabolic and Molecular Bases of Inherited Diseases. New York: McGraw-Hill 2004.
 56. Singbartl G, Walther-Wenke G, rezensiert von Frietsch T. *Transfusionspraxis – Perioperatives Management*. Berlin: Springer 2003.
 57. Singh BS, Sadiq HF, Noguchi A, Keenan WJ. Efficacy of albuterol inhalation in treatment of hyperkalemia in premature neonates. *J Pediatr* 2002; 1: 16–20.
 58. Slichter SJ. Evidence-based platelet transfusion guidelines. *Hematology Am Soc Hematol Educ Program* 2007; 172–8.
 59. Sola-Visner M, Saxonhouse MA, Brown RE. Neonatal thrombocytopenia: what we do and don't know. *Early Hum Dev* 2008; 8: 499–506.
 60. Sola-Visner M, Sallmon H, Brown R. New insights into the mechanisms of nonimmune thrombocytopenia in neonates. *Semin Perinatol* 2009; 1: 43–51.
 61. Stoffan AP, Wilson JM, Jennings RW et al. Does the ex utero intrapartum treatment to extracorporeal membrane oxygenation procedure change outcomes for high-risk patients with congenital diaphragmatic hernia? *J Pediatr Surg* 2012; 6: 1053–7.
 62. Strauss T, Sidlik-Muskatel R, Kenet G. Developmental hemostasis: primary hemostasis and evaluation of platelet function in neonates. *Semin Fetal Neonatal Med* 2011; 6: 301–4.
 63. Strohm B, Hobson A, Brocklehurst P et al. Subcutaneous fat necrosis after moderate therapeutic hypothermia in neonates. *Pediatrics* 2011; 2: e450–2.

64. Sutor AH. Vitamin K deficiency bleeding in infants and children. *Semin Thromb Hemost* 1995; 3: 317–29.
65. Thornburg C, Pipe S. Neonatal thromboembolic emergencies. *Semin Fetal Neonatal Med* 2006; 3: 198–206.
66. Turner T, Prowse CV, Prescott RJ, Cash JD. A clinical trial on the early detection and correction of haemostatic defects in selected high-risk neonates. *Br J Haematol* 1981; 1: 65–75.
67. Van Ommen CH, Heijboer H, Buller HR et al. Venous thromboembolism in childhood: a prospective two-year registry in The Netherlands. *J Pediatr* 2001; 5: 676–81.
68. Vemgal P, Ohlsson A. Interventions for non-oliguric hyperkalaemia in preterm neonates. *Cochrane Database Syst Rev* 2012; 5: CD005257.
69. Wang M, Hays T, Balasa V et al. Low-dose tissue plasminogen activator thrombolysis in children. *J Pediatr Hematol Oncol* 2003; 5: 379–86.
70. Weiss M, Schmidt J, Eich C et al. Handlungsempfehlung zur Prävention und Behandlung des unerwartet schwierigen Atemwegs in der Kinderanästhesie. *Anästh Intensivmed* 2010; Suppl 3: 54–64.
71. Weiss M, Gerber AC. Anästhesieeinleitung und Intubation beim Kind mit vollem Magen – Zeit zum Umdenken. *Anaesthesist* 2007; 56: 1210–6.
72. Wiedmeier SE, Henry E, Sola-Visner MC, Christensen RD. Platelet reference ranges for neonates, defined using data from over 47,000 patients in a multihospital healthcare system. *J Perinatol* 2009; 2: 130–6.
73. Wu YW, Hamrick SE, Miller SP et al. Intraventricular hemorrhage in term neonates caused by sinovenous thrombosis. *Ann Neurol* 2003; 1: 123–6.
74. Young G, Wicklund B, Neff P et al. Off-label use of rFVIIa in children with excessive bleeding: a consecutive study of 153 off-label uses in 139 children. *Pediatr Blood Cancer* 2009; 2: 179–83.
- Suppl 3: S876–908. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
2. Leonard E, Weisman MD, Moahn Pammi MD. Treatment and prevention of bacterial sepsis in the preterm infant. www.uptodate.com/contents/treatment-and-prevention-of-bacterial-sepsis-in-the-preterm-infant?source=search_result&search=Treatment+and+prevention+of+bacterial+sepsis&selectedTitle=2%7E150.
3. Moosmann J, Doll U, Ruffer A et al. Erste Erfahrungen mit dem Vasopressin-Rezeptor Antagonisten (Tolvaptan) als Rescue Diuretikum im Postoperativen Management bei Säuglingen mit HLHS. Stuttgart: Thieme 2013.
4. Neubert A, Botzenhardt S, Stammschulte T et al. Sildenafil zur Behandlung der pulmonalen arteriellen Hypertonie (PAH). *Monatsschr Kinderheilkd* 2013; 1029–36. Siehe hierzu auch special issue, expert consensus statement, European Pediatric Pulmonary Vascular Disease Network Heart (multipaper) 2015/2016 in press.
5. Ng E, Taddio A, Ohlsson A. Intravenous midazolam infusion for sedation of infants in the neonatal intensive care unit. *Cochrane Database Syst Rev* 2003; 1: CD002052.
6. Roehr CC, Hansmann G, Hoehn T, Bühner C. The 2010 Guidelines on Neonatal Resuscitation (AHA, ERC, ILCOR): similarities and differences – what progress has been made since 2005? *Klin Padiatr* 2011; 5: 299–307. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
7. Young T, Mangum B. *Neofax: A Manual of Drugs used in Neonatal Care*. 20th ed. Montvale, NJ: Thomson Healthcare 2010.

Teil V Notfallmedikamente zur Versorgung Neugeborener

1. Kleinman ME, Chameides L, Schexnayder SM et al. Part 14: pediatric advanced life support: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010; 18

Teil VI Transport von Früh- und Reifgeborenen

1. Beverley D, Foote K, Howel D, Congdon P. Effect of birthplace on infants with low birth weight. *Br Med J (Clin Res Ed)* 1986; 293: 981–3.
2. Das UG, Leuthner SR. Preparing the neonate for transport. *Pediatr Clin North Am* 2004; 3: 581–98, vii.
3. Deutsche Gesellschaft für Neonatologie und Pädiatrische Intensivmedizin, Deutsche Gesellschaft für Gynäkologie und Geburtshilfe,

- Deutsche Gesellschaft für Kinderheilkunde und Jugendmedizin, Deutsche Gesellschaft für Perinatale Medizin. Verlegung von Früh- und Reifgeborenen in Krankenhäuser der adäquaten Versorgungsstufe (AWMF-Register Nr. 024/002; www.awmf.org).
4. Fenton AC, Ainsworth SB, Sturgiss SN. Population-based outcomes after acute antenatal transfer. *Paediatr Perinat Epidemiol* 2002; 3: 278–85.
 5. Field D, Draper ES. Survival and place of delivery following preterm birth: 1994–96. *Arch Dis Child Fetal Neonatal Ed* 1999; 2: F111–4.
 6. McNamara PJ, Mak W, Whyte HE. Dedicated neonatal retrieval teams improve delivery room resuscitation of outborn premature infants. *J Perinatol* 2005; 5: 309–14.
 7. Mori R, Fujimura M, Shiraishi J et al. Duration of inter-facility neonatal transport and neonatal mortality: systematic review and cohort study. *Pediatr Int* 2007; 4: 452–8.
 8. Ozminkowski RJ, Wortman PM, Roloff DW. Inborn/outborn status and neonatal survival: a meta-analysis of non-randomized studies. *Stat Med* 1988; 12: 1207–21.
 9. Rüdiger M. Perinatale Asphyxie und Hypothermiebehandlung. Ligatur Verlag für Klinik und Praxis 2009.
 10. Shlossman PA, Manley JS, Sciscione AC, Colmorgen GH. An analysis of neonatal morbidity and mortality in maternal (in utero) and neonatal transports at 24–34 weeks' gestation. *Am J Perinatol* 1997; 8: 449–56.
 11. Van Reempts P, Gortner L, Milligan D et al. Characteristics of neonatal units that care for very preterm infants in Europe: results from the MOSAIC study. *Pediatrics* 2007; 4: e815–25.
 3. Deutsche Gesellschaft für Gynäkologie und Geburtshilfe, Deutsche Gesellschaft für Kinder- und Jugendmedizin, Deutsche Gesellschaft für Perinatale Medizin, Akademie für Ethik in der Medizin, Gesellschaft für Neonatologie und Pädiatrische Intensivmedizin, Deutscher Hebammenverband, Bundesverband »Das frühgeborene Kind«. Frühgeborene an der Grenze der Lebensfähigkeit (AWMF-Register Nr. 024/019; bitte Update von 04/2014 lesen; www.awmf.org).
 4. Grundgesetz der Bundesrepublik Deutschland. www.gesetze-im-internet.de/gg/BJNR000010949.html.
 5. Hintz SR, Kendrick DE, Wilson-Costello DE et al. Early-childhood neurodevelopmental outcomes are not improving for infants born at <25 weeks' gestational age. *Pediatrics* 2011; 1: 62–70.
 6. Kattwinkel J, Perlman JM, Aziz K et al. Part 15: neonatal resuscitation: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010; 18 Suppl 3: S909–19.
 7. Österreichische Gesellschaft für Kinder- und Jugendheilkunde. Erstversorgung von Frühgeborenen an der Grenze der Lebensfähigkeit. *Monatsschr Kinderheilkd* 2005: 711–15.
 8. Richmond S, Wyllie J. European Resuscitation Council Guidelines for Resuscitation 2010 Section 7. Resuscitation of babies at birth. *Resuscitation* 2010; 10: 1389–99. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
 9. Roehr CC, Hansmann G, Hoehn T, Bühner C. The 2010 Guidelines on Neonatal Resuscitation (AHA, ERC, ILCOR): similarities and differences – what progress has been made since 2005? *Klin Padiatr* 2011; 5: 299–307. Siehe hierzu die neuen internationalen ILCOR-Guidelines on neonatal resuscitation; erscheinen Ende 2015 in *Circulation and Resuscitation*.
 10. Schweizerische Akademie der Medizinischen Wissenschaften. Jahresbericht 1999.
 11. Tyson JE, Parikh NA, Langer J et al. Intensive care for extreme prematurity – moving beyond gestational age. *N Engl J Med* 2008; 16: 1672–81.
 12. Wissenschafte SAdM. Medizinisch-ethische Richtlinien zu Grenzfragen der Intensivmedizin. www.samw.ch/de/Ethik/Richtlinien/Aktuell-gueltige-Richtlinien.html.

Teil VII Entscheidungsfindung in ethischen Grenzsituationen

1. Berger TM, Bernet V, El Alama S et al. Perinatal care at the limit of viability between 22 and 26 completed weeks of gestation in Switzerland. 2011 revision of the Swiss recommendations. *Swiss Med Wkly* 2011; 141: w13280.
2. Bundesärztekammer. Grundsätze der Bundesärztekammer zur ärztlichen Sterbebegleitung. *Dtsch Ärztebl* 2011; 108: A346–8. www.bundesaeztekammer.de/downloads/Sterbebegleitung_17022011.pdf.

13. Wissenschaften SAdM. Behandlung und Betreuung von zerebral schwerst geschädigten Langzeitpatienten.
www.samw.ch/de/Ethik/Richtlinien/Aktuell-gueltige-Richtlinien.html.
2. Roos R, Proquitté H, Genzel-Boroviczény O. Checkliste Neonatologie. Stuttgart: Thieme 2010.
3. Sitzmann FC, Bartmann P, Roos R. Erkrankungen in der Neugeborenenperiode. Duale Reihe Pädiatrie. Stuttgart: Thieme 2002.

Teil VIII Anhang

1. Kattner E MB, Keen DV, Pearse RG, Dudenhausen JW. Perzentilenkurven für Geburtsgewicht, Länge und Kopfumfang unter besonderer Berücksichtigung sehr unreifer Frühgeborener. *Perinatalmed* 1992; 4: 118–21.
4. Von Versmold HT KJ, Phibbs RH, Gregory GA, Tooley WH. Aortic blood pressure during the first 12 hours of life in infants with birth weight 610 to 4220 grams. *Pediatrics* 1981; 67: 607–13.
5. Hansmann G (ed). *Neonatal Emergencies. A Practical Guide for Resuscitation, Transport and Critical Care of Newborn Babies*. Cambridge: Cambridge University Press 2009.